#### METHODS AND COMPOSITIONS FOR DIAGNOSING PALMOPLANTAR KERATODERMAS AND DYSPLASIAS AND OTHER PERIODONTAL DISEASES

5

10

15

20

25

30

35

Pursuant to 35 U.S.C. §202(c), it is acknowledged that the U.S. Government has certain rights in the invention described herein, which was made in part with funds from NIDCR, Grant Numbers: DE11601 and DE12920.

#### FIELD OF THE INVENTION

The present invention relates to the fields of genetic screening and molecular biology. specifically, the invention provides compositions and methods that may be used to advantage to isolate and detect a palmoplantar keratoderma predisposing gene, cathepsin C (CTSC), some mutant alleles of which cause susceptibility to certain pathological disorders, in particular Papillon-LeFevre Syndrome, Haim-Munk Syndrome and certain forms of early onset periodontal diseases. More specifically, the invention relates to germline mutations and functional polymorphisms in the CTSC gene and their use in the diagnosis of predisposition to palmoplantar ectodermal disorders/ dysplasias and periodontal diseases. The invention also relates to the therapy of palmoplantar ectodermal disorders/dysplasias and periodontal diseases which have a mutation or functional polymorphisms in the CTSC gene, including gene therapy, protein replacement therapy and protein mimetics. The invention further relates to the screening of drugs for treating and alleviating disease symptoms. Finally, the invention relates to the screening of the CTSC gene for disease-related mutations, which are useful for diagnosing the predisposition to additional disorders and dysplasias, including but not limited to prepubertal periodontitis, early onset periodontal disease or other forms of gum disease.

WO 01/07663

5

10

15

20

25

30

35

#### BACKGROUND OF THE INVENTION

Various publications or patents may be referenced in this application by numerals in parentheses to describe the state of the art to which the invention pertains. Full citations for these references are provided at the end of the specification. Each of these publications or patents is incorporated by reference herein.

Most forms of inflammatory periodontal disease can be successfully treated and managed. As a result, the ultimate goal of periodontal therapy has changed from that of simply arresting disease progression to one aimed at regenerating the supporting tissues.

Unfortunately, not all forms of periodontal disease respond to treatment. Severe periodontal that is resistant to conventional periodontal treatment has been recognized in a number of monogenic conditions.

Certainly some of the most intriguing and dentally challenging of these conditions include Papillon-Lefevre syndrome (PLS), Haim-Munk syndrome (HMS) and periodontal diseases.

In 1924, Papillon and Lefevre described two siblings, the products of a first cousin mating, with a condition characterized by diffuse transgradient palmoplantar keratosis (PPK) and the premature loss of both the decidous and permanent dentitions. condition came to be known as Papillon-Lefevre syndrome and subsequently over 200 cases have been described. The hallmarks of PLS are palmoplantar keratosis and rapid periodontal destruction of both dentitions. increased susceptibility to infection has been reported in approximately 20% of PLS patients. Additional findings include intracranial calcifications, retardation of the somatic development, follicular hyperkeratosis and onychogryphosis. Clinical findings reported in PLS patients suggest that the clinical expression of the condition is 5

10

15

20

25

30

PCT/US00/20400

highly variable. Unfortunately, to date, no pathognomonic disease marker exists allowing definitive diagnosis of PLS.

In 1965, Haim and Munk described an unusual syndrome in four siblings of a Jewish religious isolate from Cochin, India [21]. In addition to congenital palmoplantar keratosis and progressive early onset periodontal destruction, other clinical findings shared by these individuals included recurrent pyogenic skin infections, acroosteolysis, atrophic changes of the nails, arachnodactyly, and a peculiar radiographic deformity of the fingers consisting of tapered pointed phalangeal ends and a clawlike volar curve. Subsequently pes planus was reported to be associated with the syndrome [24]. This was the first reported association of these clinical findings, and the condition became known as Haim Munk syndrome, or keratosis palmoplantaris with periodontopathia and onychogryposis (HMS; MIM245010)[22]. Although the palmoplantar findings and severe periodontitis were suggestive of the Papillon-Lefevre syndrome (PLS; MIM245000)[3], the association of other clinical features, particularly nail deformities and arachnodactyly, argued that HMS was a distinct disorder.

PLS and HMS are classified as type IV palmoplantar ectodermal keratodermas [2]. The unique presence of severe, early onset periodontitis distinguishes PLS and HMS from other PPKs and raises the question of whether they result from the variable clinical expression of a common gene mutation, are allelic mutations at the same genetic locus, or result from expression of gene mutations at separate loci. Although Haim and Munk's initial report proposed HMS was a distinct entity, Hacham-Zadeh and co-workers referred to the disorder as Papillon-Lefevre syndrome and cited Gorlin's suggestion that HMS was a clinical variant of PLS [23,25]. In his review of PLS cases reported in the literature, Haenke

WO 01/07663

5

10

15

20

25

30

35

[5] summarizes an extensive list of clinical findings reported in PLS affected individuals, including increased susceptibility to infections, ectopic cranial calcifications and nail anomalies [5,26]. It is unclear if these additional clinical features are coincidental findings that may be segregating in a particular family or if they are etiologically related to a syndrome with a very variable clinical expression. Because PLS is an uncommon condition, and generally occurs only in a single generation it is difficult to determine if these occasional reports of associated clinical findings are etiologically related to PLS. Additionally, consanguinity is common among parents of PLS cases and therefore, it may be expected that an increased number of rare recessive conditions may be seen. Such is likely the case for the reports of mental retardation associated with PLS [5].

Pre-pubertal periodontitis (PPP) is a rare and rapidly progressive disease that results in destruction of the periodontal support of the primary dentition. The condition may be localized (usually to deciduous molars) or generalized. The localized form begins at approximately 4 years of age and is associated with only mild gingival inflammation in the presence of relatively little plaque. The generalized form begins earlier, immediately after eruption of the deciduous teeth. It is associated with severe gingival inflammation and hyperplasia, although significant gingival recession has also been described as an associated clinical feature. The attachment loss appears to be continuous rather than intermittent as with most other forms of periodontitis.

A varied clinical phenotype has been reported for PPP, probably reflecting the fact that the term PP describes an etiologically heterogeneous group of conditions that share an overlapping clinical presentation. Although PPP can occur as an isolated finding, many reports of PPP describe an increased

systemic susceptibility to infections. Children with PPP have frequent inner ear infections and infections of the respiratory tract [39,40]. Prepubertal periodontitis is known to be associated with Papillon Lefevre syndrome and with a number of systemic disease states that share an increased susceptibility to microbial infections.

To date, no pathognomonic disease marker exists for most PPKs allowing for definitive diagnosis. The present invention provides such a disease marker and methods of use thereof having diagnostic and prognostic utilities for several PPKs and many periodontal diseases.

#### SUMMARY OF THE INVENTION

5

10

15

20

25

30

35

The present invention provides compositions and methods which allow for genetic screening and diagnosis of certain palmoplantar keratodermas and periodontal disease states in affected individuals. In accordance with the present invention, it has been discovered that mutations or functional polymorphisms in the cathepsin C gene (CTSC) give rise to certain pathological conditions including PLS, HMS and periodontal diseases. Mutations or functional polymorphisms associated with the disease state are those which give rise to a altered, truncated, misfolded or otherwise non-functional CTSC polypeptides. Polymorphisms in the CTSC sequence which do not affect the nature of the encoded protein are not associated with PLS, HMS or periodontal disease.

Thus, in one embodiment of the invention, a method is provided for determining the presence of alterations in CTSC encoding nucleic acids which give rise to altered CTSC proteins. The wild-type CTSC nucleic acid sequence and its encoded amino acid sequence are known. See SEQ ID NOS: 1-3 provided herein. This sequence information facilitates the identification of genetic changes that give rise to aberrant CTSC proteins.

CTSC mutations specifically associated with PLS,

PCT/US00/20400

WO 01/07663

HMS and PPP are described herein and are set forth in Table 1. Accordingly in one embodiment of the invention, nucleic acid molecules encoding altered CTSC proteins are considered to be within the scope of the present invention. In a preferred embodiment of the invention, the altered CTSC nucleic acid has at least one of the alterations set forth in Table 1.

In a further embodiment of the invention, nucleic acid probes which specifically hybridize to the human altered CTSC-encoding nucleic acids described herein and not to wild-type CTSC encoding nucleic acids are provided. In a preferred embodiment, the probes specifically hybridize with altered CTSC encoding nucleic acids having at least one of the alterations set forth in Table 1.

In yet another embodiment of the invention, a mutated CTSC protein encoded by the altered CTSC encoding nucleic acids of the invention is provided. Preferably such CTSC proteins are encoded by a nucleic acid containing a mutation as set forth in Table 1. Also provided are assays for biochemically assessing altered cathepsin C activity. Antibodies immunologically specific for altered CTSC proteins are also contemplated to be within the scope of the present invention.

In another aspect of the invention, a method for detecting a germline alteration in a CTSC gene is provided. In a preferred embodiment the alteration is selected from the group consisting of the alterations set forth in Table 1. The method comprises analyzing a sequence of a CTSC gene or CTSC RNA from a human sample or analyzing a sequence of CTSC cDNA made from mRNA from a human sample and comparing sequences so isolated to the wild type sequence encoding CTSC. Inasmuch as certain alterations of the CTSC coding sequence may not alter the function of CTSC, methods are provided for assessing the enzymatic activity of proteins encoded by nucleic acid molecules which do not possess the wild

35

30

5

10

15

20

WO 01/07663

5

10

15

20

25

30

35

type CTSC sequence.

In yet another embodiment of the invention, kits are provided for detecting the presence of an altered CTSC encoding nucleic acids in a biological sample. exemplary kit comprises the following: oligonucleotides which specifically hybridize with CTSC encoding nucleic acids having the alterations set forth in Table 1; ii) reaction buffer; and iii) instruction sheet. Kits for detecting the presence an altered CTSC proteins in a biological sample are also Exemplary kits for this purpose comprise: i) provided. antibodies immunologically specific for the altered CTSC ii) a solid support with proteins of the invention; immobilized CTSC antigens as a positive control; and an instruction sheet. Optionally, anti-CTSC antibodies used for this purpose may contain a detectable label or tag for used in isolating or detecting immune complexes.

Various terms relating to the biological molecules and cells of the present invention are used throughout the specifications and claims.

With reference to nucleic acids used in the invention, the term "isolated nucleic acid" is sometimes employed. This term, when applied to DNA, refers to a DNA molecule that is separated from sequences with which it is immediately contiguous (in the 5' and 3' directions) in the naturally occurring genome of the organism from which it was derived. For example, the "isolated nucleic acid" may comprise a DNA molecule inserted into a vector, such as a plasmid or virus vector, or integrated into the genomic DNA of a procaryote or eucaryote. An "isolated nucleic acid molecule" may also comprise a cDNA molecule. isolated nucleic acid molecule inserted into a vector is also sometimes referred to herein as a Arecombinant@ nucleic acid molecule.

With respect to RNA molecules, the term "isolated

nucleic acid" primarily refers to an RNA molecule encoded by an isolated DNA molecule as defined above. Alternatively, the term may refer to an RNA molecule that has been sufficiently separated from RNA molecules with which it would be associated in its natural state (i.e., in cells or tissues), such that it exists in a "substantially pure" form (the term "substantially pure" is defined below).

With respect to protein, the term "isolated protein" or "isolated and purified protein" is sometimes This term refers primarily to a protein used herein. produced by expression of an isolated nucleic acid molecule of the invention. Alternatively, this term may refer to a protein which has been sufficiently separated from other proteins with which it would naturally be associated, so as to exist in "substantially pure" form.

The term "substantially pure" refers to a preparation comprising at least 50-60% by weight the compound of interest (e.g., nucleic acid, oligonucleotide, protein, etc.). More preferably, the preparation comprises at least 75% by weight, and most preferably 90-99% by weight, the compound of interest. Purity is measured by methods appropriate for the compound of interest (e.g. chromatographic methods, agarose or polyacrylamide gel electrophoresis, HPLC analysis, and the like).

With respect to antibodies, the term "immunologically specific" refers to antibodies that bind to one or more epitopes of a protein of interest, but which do not substantially recognize and bind other molecules in a sample containing a mixed population of antigenic biological molecules.

With respect to single stranded nucleic acids, particularly oligonucleotides, the term "specifically hybridizing" refers to the association between two single-stranded nucleotide molecules of sufficiently complementary sequence to permit such hybridization

35

30

5

10

15

20

under pre-determined conditions generally used in the art (sometimes termed "substantially complementary"). In particular, the term refers to hybridization of an oligonucleotide with a substantially complementary sequence contained within a single-stranded DNA or RNA molecule of the invention, to the substantial exclusion of hybridization of the oligonucleotide with single-stranded nucleic acids of non-complementary sequence. Appropriate conditions enabling specific hybridization of single stranded nucleic acid molecules of varying complementarity are well known in the art.

For instance, one common formula for calculating the stringency conditions required to achieve hybridization between nucleic acid molecules of a specified sequence homology is set forth below (Sambrook et al., 1989):

 $T_m = 81.5$ °C + 16.6Log [Na+] + 0.41(% G+C) - 0.63 (% formamide) - 600/#bp in duplex

20

25

30

35

5

10

15

As an illustration of the above formula, using [Na+] = [0.368] and 50% formamide, with GC content of 42% and an average probe size of 200 bases, the  $T_m$  is 57°C. The  $T_m$  of a DNA duplex decreases by 1 - 1.5°C with every 1% decrease in homology. Thus, targets with greater than about 75% sequence identity would be observed using a hybridization temperature of 42°C.

The term "promoter region" refers to the transcriptional regulatory regions of a gene, which may be found at the 5' or 3' side of the coding region, or within the coding region, or within introns. In the present invention, the use of SV40, TK, Albumin, SP6, T7 gene promoters, among others, is contemplated. Specific promoters for the yeast and mammalian expression systems of the invention are available and known to those of ordinary skill in the art.

The term "operably linked" means that the

regulatory sequences necessary for expression of the coding sequence are placed in the DNA molecule in the appropriate positions relative to the coding sequence so as to enable expression of the coding sequence. This same definition is sometimes applied to the arrangement of transcription units and other transcription control elements (e.g. enhancers) in an expression vector.

5

10

15

20

25

30

35

The phrase "functional polymorphism" refers to a change in wild type CTSC coding sequence giving rise to altered cathepsin C activity as assayed using conventional methods.

The term "oligonucleotide," as used herein refers to primers and probes of the present invention, and is defined as a nucleic acid molecule comprised of two or more ribo- or deoxyribonucleotides, preferably more than three. The exact size of the oligonucleotide will depend on various factors and on the particular application and use of the oligonucleotide.

The term "probe" as used herein refers to an oligonucleotide, polynucleotide or nucleic acid, either RNA or DNA, whether occurring naturally as in a purified restriction enzyme digest or produced synthetically, which is capable of annealing with or specifically hybridizing to a nucleic acid with sequences complementary to the probe. A probe may be either single-stranded or double-stranded. The exact length of the probe will depend upon many factors, including temperature, source of probe and use of the method. example, for diagnostic applications, depending on the complexity of the target sequence, the oligonucleotide probe typically contains 15-25 or more nucleotides, although it may contain fewer nucleotides. herein are selected to be complementary to different strands of a particular target nucleic acid sequence. This means that the probes must be sufficiently complementary so as to be able to "specifically hybridize" or anneal with their respective target

IC

strands under a set of pre-determined conditions. Therefore, the probe sequence need not reflect the exact complementary sequence of the target. For example, a non-complementary nucleotide fragment may be attached to the 5' or 3' end of the probe, with the remainder of the probe sequence being complementary to the target strand. Alternatively, non-complementary bases or longer sequences can be interspersed into the probe, provided that the probe sequence has sufficient complementarity with the sequence of the target nucleic acid to anneal therewith specifically.

5

10

15

20

25

30

35

The term "primer" as used herein refers to an oligonucleotide, either RNA or DNA, either single-stranded or double-stranded, either derived from a biological system, generated by restriction enzyme digestion, or produced synthetically which, when placed in the proper environment, is able to functionally act as an initiator of template-dependent nucleic acid synthesis. When presented with an appropriate nucleic acid template, suitable nucleoside triphosphate precursors of nucleic acids, a polymerase enzyme, suitable cofactors and conditions such as a suitable temperature and pH, the primer may be extended at its 3' terminus by the addition of nucleotides by the action of a polymerase or similar activity to yield an primer extension product. The primer may vary in length depending on the particular conditions and requirement of the application. For example, in diagnostic applications, the oligonucleotide primer is typically 15-25 or more nucleotides in length. The primer must be of sufficient complementarity to the desired template to prime the synthesis of the desired extension product, that is, to be able anneal with the desired template strand in a manner sufficient to provide the 3' hydroxyl moiety of the primer in appropriate juxtaposition for use in the initiation of synthesis by a polymerase or similar enzyme. It is not required that the primer

sequence represent an exact complement of the desired template. For example, a non-complementary nucleotide sequence may be attached to the 5' end of an otherwise complementary primer. Alternatively, non-complementary bases may be interspersed within the oligonucleotide primer sequence, provided that the primer sequence has sufficient complementarity with the sequence of the desired template strand to functionally provide a template-primer complex for the synthesis of the extension product.

## BRIEF DESCRIPTION OF THE DRAWINGS

Figures 1A-1E are a series of clinical photographs showing palmoplantar keratosis and periodontal disease in PLS study patient. Fig. 1A: palmar hyperkeratotic lesions; Fig. 1B: plantar hyperkeratotic lesions; Fig. 1C: hyperkeratotic lesions affecting the knees; Fig. 1D: periodontitis involving erupting permanent dentition and Fig. 1E: periapical radiographs showing severe alveolar bone loss affecting erupting permanent teeth.

Figure 2 shows haplotype data for chromosome 11q short tandem repeat polymorphisms (STRP) markers spanning the PLS gene locus. Segments which are likely to be homozygous by descent are boxed. Arrows indicate recombinant events. Individuals 7 and 22 share a common haplotype for D11S1979, D11S1887, D11S1780, D11S1367, D11S931, and D11S4175.

30

5

10

15

20

25

Figure 3 depicts pedigree and sequence analysis of CTSC exon 6. The numbering of the wildtype sequence shown above the figure is based upon the genomic sequence of CTSC. See SEQ ID NO: 1. Circles represent females and squares represent males. Filled symbols indicate affected individuals. Half-shading indicates carriers based upon DNA sequencing results. All affected individuals are homozygous for the specific CTSC

mutations. Arrows indicate the position of the mutation. This family has a nonsense mutation (856 C->T) at codon 286 resulting in a truncated protein of 286 amino acids.

5

10

15

20

25

30

35

Figures 4A - 4D show pedigrees and sequence analysis of CTSC exon 7 for 4 Families with PLS. Symbols are as described for Figure 3. Fig. 4A: Family with a single base pair deletion (1047delA) of CTSC resulting in a frameshift and premature termination. Fig. 4B: Family with a 2bp deletion (1028-1029delCT) of CTSC resulting in a frameshift and premature termination. Fig. 4C. and Fig. 4D: pedigrees of families with a nonsense mutation (1286G->A) at codon 429 resulting in a truncated protein of 428 amino acids. The father in family C is deceased and no sample was available for analysis.

Figures 5A-5C is a schematic diagram of the CTSC gene showing the locations of the mutations described herein. Panel 5A. Genomic structure of CTSC gene with introns shown as solid lines and exons depicted as boxes. The 5' and 3' untranslated regions are shown as filled boxes. Panel 5B. Coding region of CTSC gene. The amino acid numbers are shown at the end of each exon. Mutations listed in Table 1 are shown according to their genomic locations with Missense, ★; Nonsense, ♦; Insertion, ■; and Deletion,□. The splicing site mutation is indicated by an arrow. Panel 5C. Subunit structure of CTSC polypeptide with SP, signal peptide; P1, 13.5 kDa pro-region; P2, 10 kDa pro-region; H, heavy chain; and L, light chain. The 10 kDa pro-region is cleaved out upon activation. The disulfide bond within the 13.5 pro-region is shown. The glycosylation sites are indicated by filled circle and arrows indicate the active sites.

Figures 6A-6F show a series of micrographs

depicting the clinical and radiographic findings in Haim Munk syndrome. Fig. 6A: dermal involvement of fingers in individual #34. Fig. 6B: Individual #34 radiograph of terminal phalanges of the fingers showing marked thinning increasing towards the distal, tapering pointed ends showing a claw-like volar bend. Fig. 6C: Individual #17 palmar keratosis; Fig. 6D: Individual #17 plantar keratosis: Fig. 6E: Individual #17 gingival inflammation; Fig. 6F: Individual #17 radiograph showing alveolar bone destruction associated with gingival inflammation shown in 6D.

Figure 7A shows pedigree of Cochin descendents segregating Haim Munk syndrome (HMS). Numbered individuals have been analyzed for the current study. Circles = females, squares = males, shaded symbols = HMS affected individuals. Double lines = consanguinity. Individuals #10,11 \* = second cousins. Numbers inside circles, squares and diamonds indicate the number of additional offspring not examined in this study. Sibships described in previous reports are indicated and referenced below the pedigrees. The subjects of Haim and Munks original report (1965) are individuals 33, 34, 35, and 36. Half-shading indicates carriers based upon DNA sequencing and/or restriction enzyme analysis. Unshaded numbered individuals represent non-carriers based upon DNA analysis. Figure 7B shows a pedigree of a Turkish family segregating PLS. Numbered individuals were available for study. Half-shading indicates carrier based upon DNA analysis. Individual 77 is a non-carrier based upon DNA sequencing.

Figures 8A and 8B show the results of sequence analysis of exon 6 of CTSC. The numbering of the wildtype sequence is based upon the cDNA sequence of CTSC. See SEQ ID NO: 1. Fig. 8A: Family A (Cochin isolate diagnosed with Haim Munk syndrome) from Figure

35

30

5

10

15

20

Affected individuals are homozygous for a 857A->G missense mutation which results in a conserved glutamine being changed to an arginine (Q286R). Representative sequences are shown for individuals #36 (affected) and #31(carrier). Fig. 8B. Family B from Figure 1. Affected individuals are homozygous for a 856C->T nonsense mutation which results in a premature stop codon at position 286 (Q286X). The Q286X mutation has been previously reported in an inbred Turkish family [12].

5

10

15

20

25

30

35

Figure 9 depicts a gel showing the results of restriction enzyme analysis of Q286R mutation in Family A of Fig. 7. A 465 bp fragment of exon 6 was amplified and subjected to restriction digestion with AvaI as described under methods. The Q286R mutation introduces a new Aval site. After digestion and electrophoresis through 1.8% agarose gels, wildtype individuals exhibit bands of 465 bp, affected individuals have bands of 404 and 61 bp, and carriers have bands of 465, 404, and 61 M. 1 kb ladder (Gibco). Lane 1. Individual #5 uncut, demonstrating 465bp amplicon. Lane 2. Individual #5 cut with AvaI. Only the 465 bp fragment is observed. Thus individual #5 has the wildtype sequence on both alleles. Lane 3. Individual #31 uncut. Lane 4. The 465 and 404bp Individual # 31 cut with AvaI. fragments are visible, confirming that individual #31 is a carrier of the Q286R, consistent with the sequencing results shown in Figure 3A. Lane 5. Individual #34 uncut. Lane 6. Individual #34 cut with AvaI. The 404 and 61bp fragments are indicated by arrows.

# DETAILED DESCRIPTION OF THE INVENTION

The present invention relates generally to the field of human genetics. Specifically, the present invention relates to methods and materials used to isolate and detect mutated forms of the lysosomal

protease cathepsin C (CTSC) gene, associated with autosomal recessive disorders characterized by palmar hyperkeratosis and/or periodontitis. More specifically, the present invention relates to germline mutations in the CTSC gene and their use in the diagnosis of predisposition to such pathological conditions. Additionally, the invention relates to germline mutations in the CTSC gene in other palmoplantar ectodermal disorders and dysplasias and their use in the diagnosis and prognosis of such pathological conditions. The invention also relates to the therapy of palmoplantar ectodermal disorders and dysplasias which have a mutation or functional polymorphism in the CTSC gene, including gene therapy, protein replacement therapy and protein mimetics. The invention further relates to the screening of drugs which may have therapeutic value. Biochemical assays are provided for the assessment of altered activity of aberrant CTSC enzymes encoded by the mutated CTSC encoding nucleic acids of the invention. Finally, the invention relates to the screening of the CTSC gene for mutations, which are useful for diagnosing the predisposition to ectodermal disorders and dysplasias.

The present invention provides an isolated polynucleotide comprising all, or a portion of the CTSC locus or of a mutated CTSC locus, preferably at least eight bases and not more than about 100 kb in length. Such polynucleotides may be antisense polynucleotides. The present invention also provides a recombinant construct comprising such an isolated polynucleotide, for example, a recombinant construct suitable for expression in a transformed host cell.

Also provided by the present invention are methods of detecting a polynucleotide comprising a portion of the CTSC locus or its expression product in an analyte. Such methods may further comprise the step of amplifying the portion of the CTSC locus, and may further include a

35

30

5

10

15

20

5

10

15

20

25

step of providing a set of polynucleotides which are primers for amplification of said portion of the CTSC locus. The method is useful for either diagnosis of the predisposition to PPKs or the diagnosis or prognosis of keratodermal disorders/dysplasias and periodontal diseases.

The present invention also provides isolated antibodies, preferably monoclonal antibodies, which specifically bind to an isolated polypeptide comprised of at least five amino acid residues encoded by the altered CTSC locus.

The present invention also provides kits for detecting in an analyte a polynucleotide comprising a portion of the CTSC locus, the kits comprising a polynucleotide complementary to the portion of the CTSC locus packaged in a suitable container, and instructions for its use.

The present invention further provides methods of preparing a polynucleotide comprising polymerizing nucleotides to yield a sequence comprised of at least eight consecutive nucleotides of the CTSC locus; and methods of preparing a polypeptide comprising polymerizing amino acids to yield a sequence comprising at least five amino acids encoded within the CTSC locus.

The present invention further provides methods of screening the CTSC gene to identify mutations. Such methods may further comprise the step of amplifying a portion of the CTSC locus, and may further include a step of providing a set of polynucleotides which are primers for amplification of said portion of the CTSC locus. Exemplary primers are set forth in Table A.

The method is useful for identifying mutations for use in either diagnosis of the predisposition to keratodermal disorders/dysplasias and periodontal diseases or the diagnosis of such disorders.

The present invention further provides methods of screening suspected CTSC mutant alleles and functional

35

polymorphisms to identify mutations in the CTSC gene. In addition, the present invention provides methods of screening drugs for therapy and to identify suitable drugs for restoring CTSC gene product function.

Finally, the present invention provides the means necessary for production of gene-based therapies directed at aberrant cells associated with keratodermal disorders and dysplasias. These therapeutic agents may take the form of polynucleotides comprising all or a portion of the CTSC locus placed in appropriate vectors or delivered to target cells in more direct ways such that the function of the CTSC protein is reconstituted. Therapeutic agents may also take the form of polypeptides based on either a portion of, or the entire protein sequence of CTSC. These may functionally replace the activity of CTSC in vivo.

It is a discovery of the present invention that mutations in the CTSC locus in the germline are indicative of a predisposition to keratodermal disorders/dysplasias and periodontal diseases. The mutational events of the CTSC locus can involve deletions, insertions and point mutations within the coding sequence and the non-coding sequence.

A major gene locus associated with the keratodermal disorders and dysplasias of the invention has been localized to a 2.8 cM interval on chromosome 11q14 of the human genome. This region contains a genetic locus, The CTSC message is expressed at high levels in a variety of immune cells including polymorphonuclear leukocytes, macrophages and their precursors. This gene is expressed in the palms, soles, knees, and oral keratinized gingiva.

The CTSC gene was originally reported to consist of 2 exons. US Provisional Application 60/165,016 from which the present application claims priority, describes mutations in Exons 1 and 2. The mutations described actually fall within Exons 6 and 7. Reference numerals

35

30

5

10

15

20

to the altered amino acids are the same as those in US Provisional 60/165,016, only the nucleotide numbering has changed to reflect the actual genomic structure of the CTSC gene which is now known to contain 7 exons. The sequence encoding the wild type human CTSC gene is provided below (SEQ ID NO: 1):

5

AGGGAGATAT AAGTGAATAA TTTGGACCTG CTCTCTTTGA ATGTTTATAA TCTGGTGGAA AAAAAATGGA CATATGAATA TTGATTTGTG ACCAGTGCAA AGGGGGCAAA AATTCATATC CCAAAGAAAA CGGGGACACA TCAGGTCTGT CTTGTTCATC ACTGTGTCCA CAGGGCCTGA 10 CACCTAGTAG GCTCAGTGGG AGAAAGGAGC CCCAATTACC AACAAAAGCC AGGAAAGAAC GGGAGGCTCT TACGGAAAAG GGTGATACTT AAACTGAGCA AGGAGGCACC TGGAAATAGT GCCACCTAAT AATTTTTGGC GATCAGACTG ACACACTAGA ACGGTTCATA AGACCAGCCT TCTCCCATTG GCTAGCTTCC TTCCTCACCC TTCTCACCCT GGGCAAGCCG CTTCCTCTCT CTGGGCCTCT TGCTTTTCCT CTGTAACATA AAAGGGGTTG AGCAATATCA TCTCTGAGAG 15 CGCCATGTGT GTGCGTGCCA GAGGGAAAAC CCCCACAACG CTAATACATC AAAACTGCAG GTTTGCACAA AAACTGAATT CTGCTGAATG CAAACAGGCA AACAGCATTT ACCAGGAAAC AAAACAAAAT CAAGCACATA AAAAAGTAGG AAGAGTTGGA AAACGGAAGG AAGATAAGTT CTCAAACAGC TGGAATAGTT GATGTTAGCT AGCGAAGTTT TTCAGAGGAA AAAACAAGAA GTTGGTTATG AGGCAAGTGG ACCTGAGAAA AAAGACTAAA GGGGAAGAAT AGCAAGTAAA 20 ACAGAACTCC ACTTGCTAGA TCTCTCCCTC TGTCGCGCTC TTTCACCTGA CCCACTCCCT TATTCCCCCC ACACCCTTTC CTTCTCCCC TACGTTACCG CACAGGAACG AAGTCTGGGT CATGTGCGGA CCGCTTGTGG CTCTTAAATC CTCTTTTTGT CACCCTGGCC GTGCAAAATT TTGAAACGTC CCTCGGCAAA AAAAATAAAA ATAAAAAAAA AAAATCTGTC CCTGGCCTCT TCCCTAGTTC TGGGTCCAGT TGCAGCCAAG TGAGGGGCAG CGCGCGCTCC CAAGTCCCCG 25 TTTCAGAGAC GCGCACGCGC CTGGCGCCCA ACCCCCAATC CCCTGCTGCT CAGTGACCCC GCCCACGGGT TTCCGGGCCG GCGTAGCTAT TTCAAGGCGC GCGCCTCGTG GTGGACTCAC CGCTAGCCCG CAGCGCTCGG CTTCCTGGTA ATTCTTCACC TCTTTTCTCA GCTCCCTGCA GCATGGGTGC TGGGCCCTCC TTGCTGCTCG CCGCCCTCCT GCTGCTTCTC TCCGGCGACG GCGCCGTGCG CTGCGACACA CCTGCCAACT GCACCTATCT TGACCTGCTG GGCACCTGGG 30 TCTTCCAGGT GGGCTCCAGC GGTTCCCAGC GCGATGTCAA CTGCTCGGTT ATGGGTAAGC CGCCGGCTCG GCAGTCCTCC GGGTCGTCCT TTCTGCCCTT GAGCCCCTAA CGCAGCGCCA CGCCAACTAC CGCTTCCCCC CAGGCAGACG CTTGTGGGTG GCCAGAGCAT CTTGACTGGA TTCGGGGACC CTTGGGGACC TTCTTCCCCG CCAGGCTCGC GAAGTTAAAG TTCATCTGCT GAGAACTTCT AACTCCACAC TTTCTTGGTT ATCTTGGGGA CTCAACACTT TGATCAAGAA 35 CTTTTTTATT CCTCCCGCTT AATTTTGTTT GCTTTGAGAG AGACTTGGGA ACTGCAATCG TTTGGTTCTC CAGTCCGATC TGGTAGCGTT ATTTTTAAAA TTTATTTTTA TTTTTTATTA CTATTTTACT AGTGAAGATA GATGAGCTCA GAGACTCTCG AGGATATAGC ATGAAGTTTT CTCTTTTTGT TAGATGGTGG GAAAGGGACT TTCTGCCCAG CGATTTTGGT TTGAGCGGGT GTTGATGAGT ACTAGAAAAC GGCTAGTACC ACTCTGCATT GTTTCATGCA TTGCAAGGAG 40 GTAAAAATTT TTTAAAAAAT TAATAACAAA GAAAACTTAA CTCTGAACCT AGTAATTAGA AATGCCCAGA GTCTGCACAA TGTTTGGCTC ATGGAAGGCT CTCAATAAAT ACCTAGTGTT TGAACATACT GGAGATATTC CATATGCCTT CAGATAACAT GGTTACCCCT AGAACAAAGA  ${}^{\prime}{}^{$  PCT/US00/20400

	ACCTAGTGAA GGGGGTGGGG TGGAAAAAAG ACAATCACTA GTTAGAAGAG TCACACTGTG
	GTCTACCCAA ACCCTCTTAC AGCCTGTGAC TTCTGCAGAG TCAGTAAAAA ATCAGCTATA
	ATTTTCTTGT CAACAGAACA AAGATTAAGT CATTTGTTAC TAAAGAAATA CCTTTTTAAC
	TGACTGTATA GCATTTCATA ATCCTGAACA TTGTAACTTT TTTTTTTTT TTTTTGAGAA
_	TGGAGTCTTG CTCTGTCACC CAGGCTGTAG TTCAGTGGCG GGATCTCGGC TCACTGCAAC
5	CTCCGCCTCC CGGGTTTAAG CGATTCTCCT GCCTCAGTCT CCCGAGTAGC TAGGATTACG
	GGCACATGCC ACCACGCCCG GCTAATATTT TGTATTTTTA GTAGAGACGG GTTTCACTGT
	GGCACATGCC ACCACGCCCC GOTTATATATATATATATATATATATATATATATATATAT
	TACTGGGATT ACAGGCGGGA GTCACCACGC CTGGCCTTTT TTTTTTTTTT
1.0	TACTGGGATT ACAGGGGGGC AGAAAGAGAT CAGGATTTGC ACTGCCCTGT CACATGCAAT
10	CTCCCATGTC AGAGCATTAC TCTCAAACAT GGAAAAACTT TAAAATACAC AACTCTCCAG
	ATGGACACAG CTGAATCATT TTCAGGAAGG CTCTGCCTAT AAAATTACCT TTGGCCTTAA
	TTCAGTAATT AAAGGCACCC ACAGGTCTGA GCCCTCATCT ATGAAGGTTA AACGTATCAT
	TTCAGTAATT AAAGGCACCC ACAGGTGTOT DOUBLE TOTATCAGC ACTAATTGGT TTTATAGGAT ACAGACCTCT AGTTTGCTGA ATAAACTTTG
	GAAGGATTCA GATTATGAGG TTCTAAACTG TAGAGCCTTT GAGGAGAAAG GTACCCCATT
15	GAAGGATTCA GATTATGAGG TECTAMETO THEOTOT AGCACATTGT CTTCCTTAGG TCTTCTCCGA ATAGATCATT TTGTGTCTTC TCCTGGGCTT AGCACATTGT CTTCCTTAGG
	TCTTCTCCGA ATAGATCATT TIGIGICITE TESTSOCIAL TOTAL TOTAL CONTROLL ATCTGATAGT CTGTTATTT ATCTTTTTTTTTTTTTTT
	ATCTGATAGT CTGTTATTT ATCTTTTGT TOTGTATAGE TGTAATTTTC AACATTCCCC
	TTTTGACTAG CATTTTGCCT TTTCTCCGTT TCTCGATAGG DDDD
	ATTTCTCCTT TTGTTCAGTG GAAAAATTTC TTCAGTGTTG GGATGGTGTG GTAGTTTTGG GGGTCTCCCT GCCTGCTGGG CTGCATAGCA TATCTCTCTT TTTTGAGACA
20	GTAGTTTTGG GGGTCTCCCT GCCTGCTGGG CTGCATAGCA TATOTOGOA AGGTCTCACT GTGTTGCCCA GGTTGGAATG CAGTTGTGCA ATCATAGCTC ACTGCAGCCT
	AGGTCTCACT GTGTTGCCCA GGTTGGAATG CAGTTGTGCA MCCATCAAA TAGCTGGGAC CCAACTCCTG GGCTCAAGCA ATCCTTCCAC CTCAGCTTCA GCCGCTCAAA TAGCTGGGAC
	CCAACTCCTG GGCTCAAGCA ATCCTTCCAC CTCAGGTTCH GGGGGTTCH GGGGCAGGGGTACAGGCAAT GCACCACCAT AACCAGCGAA TTTTTAAAAA TTATTTTTGT AGAGACAGGG
	TACAGGCAAT GCACCACCAT AACCAGCGAA TITITAAAAA TITITAAAAAA TITITAAAAAAA TITITAAAAAA TITITAAAAAA TITITAAAAAA TITITAAAAAAA TITITAAAAAAA TITITAAAAAAAA
	TCTCACTATG TTGCCCAAAC TAGTCTTGAA GTTCTGACCT CAGGGCG CAGGCCTAACTTC ATTACAGGCG TGAGCCACTG CTTCTGCTTG CATTATCTCA AATTTTTAGA GCCTAACTTC
25	ATTACAGGCG TGAGCCACTG CTTCTGCTTG CATTACTCA ATTACTCA ATTAC
	ACAGTCTTGC CTCGGAGCAG GAACTGTTTG AGGCAACAAG ATGAAAATTGA ACTAGGCAGA CTGTGCTCAA ACTGGGTAGC ATGAAAGGAA AGCAGCACAA TTAAAATTGA
	ACTAGGCAGA CTGTGCTCAA ACTGGGTAGC ATGAAAGGAA ACGTGGTGCTTTTTTT  AATTGGGGGA TTCTTTCCCC TATTGAAATC CAAATAATTT TCCTTTATTG TGCTTTTTTT
	AATTGGGGGA TTCTTTCCCC TATTGAAATC CAAATAATTT TCCTTTAGAAG CTGGATACAG
	TTTTTTCTTT AGGACCACAA GAAAAAAAAG TAGTGGTGTA CCTTCAGAAG CTGGATACAG CATATGATGA CCTTGGCAAT TCTGGCCATT TCACCATCAT TTACAACCAA GGCTTTGAGA
30	CATATGATGA CCTTGGCAAT TCTGGCCATT TCACCATCAT TTACTTTGTT GGAAGTTGGA
	TTGTGTTGAA TGACTACAAG TGGTTTGCCT TTTTTAAGGT TAGTTTTGTT GGAAGTTGGA
	TTTACATTTT CAATGATTTG ATATCTGAAA CCTCTTCTGA TTAGTAGACC CTCAGAATTT
	TAATTTTAGA TTAAGAAGAT GACCGGAATT GACACCACTC TTCCCAAGAG TAGTAGTAGT
	TGGTATAATT TTGCCACTTT ATTTAAAATT ATGATTTTAG TAGGTTTACA AAATACCAGT
35	GCTACATTTG AATATGTATA AATTATGTTT AAAATTTACA TTTTGGTAAG TATGACTAAA
	TTCTTAATTT ATTTTCCTTA TTACTACCAC TTTATTTCTA AATGTTGCCA TAGTCATTTG
	GCTTTGTTCT AAATCTGTAG GAAAGATAGA GAGATTACAC ATTTTGTTTT CTTGCAGTTA
	CTATGCTGTC CTTCCTATCA CTACCTGTTG GCTGAGGTAG TGATAGGCCT AAATGATTCA
	TTATCTTAAA TGTACTAAAT ATGTTGAGTA ATTTTTTCTT CTAAACTAAC AGAAAGAGAG
40	AACCTAGGAG TTACTCCCTT AGGCTGGTTA AAGTGAAAGG TAGCCAAGTC AACCCAGCTT
	GTTTCCTTCT CTCATTAGGA AAGAACTATT GTTCATTCTC ATAACACACT TTTTCCAATT
	GCAAACATAC TCAGGGTTAA AATAGTTTAG CACAAATTGC AGCCCATTTC ATTTGTTCTT
	CACAAGCTGG AACTTTTCTT GTAAGCTAAA TATTAAATGG TTCAAGTAAA TTGGATACAT
	AAGCCTGAAA CTAGGCGTTT CTCATTATAC ATAGAGTATA AATTAAGACA GACTTTTTCA

WO 01/07663

	TCCTCAAAGG	TTTACAGCCT	ттаааасатс	TGGGAAGAAG	TGGGAAAGTA	GGGAATAACT
		TGATAAAAGA				
		AGTTGTGGTT				
		ATAACTCCCA				
5		GAAATAGAGA				
J		GTGGAAACCA				
		ACCCATGCTT				
		TTTGTAAAAA				
		TGGTATATTT				
10		GACCTCTTAC				
10		CAATTGCATG				
		ССАТАААССТ				
		TTTGACACAC				
		ATTCAGTTGG				
15		CAAGTACATA				
		GGAGAAACTG				
		TAGACAAAAG				
		GTAGTCCAAA				
		CACTTTTTGTG				
20	AGAACATTAG	G TCACATTAGE	TCAATAGAAA	ATCACTTTTT	AAAGCCAAAT	AAAGTTATAT
	GTGTTCCCA	A CCATGAGTTO	GAAAGAATTA	ATATATATGC	TGTTGGAGGG	TAGAACCCTG
	CCTAATCAT	A TGGTTCTGG	TGGCATTGAT	CGAATCCTTA	TTCTTTCATT	AGGAATAACA
	ATAGAAAAA	A TACTCCTGCC	CTACTGATT	CAGGATATGI	СТАТТТТААА	GTGCCCATTT
	GACAAAACC	A TTATCAGGG	CATGTTTTC	TTTTCTGCAG	AAAAATCAAC	CACTCTGGTC
25	AGTAGTTAG	G TCTTATGAC	A AGCACCATA	A TTTCCTTAGG	CAGAGTAGAA	TATAATAGGA
	TACTTCTTT	T TGAAACTTA	A TATAATCAG	G TAGTTCCAGA	TAAACATAGO	TTGCAAAGTG
		C ATGTTATTT				
	AAACTTAAG	A AAGATATTA	A GATGGAGTT	G ACTTTGAATA	ATAAAGTCAI	CCACTGTTGA
		ATAATTAATT				
30		G GGAAACCGT				
						TTACATTAAA
						TTCAAAGAGT
						ACAAATTCTT
						TCACGCCTCT
35						G CTTCAGATAT
						r AGTTGGTACA
•						r GGCTCATAAC
						r CTCCTACCCT
						C CCTCATACCA
40						CAAGAAGAAT
						г ататтттаас
						G CCCAAGAAGA
						A GGTGAAGAAG
	AACTCATC	AT GCTGGAAG	GG TGGTCCAC(	CC TAACTCCGC 2/	A GGGACAGAA	G CTCCTGTGTC

	TGGGACCCTT	CCAGAACTTG	CTCTATGTAT	CTCTTCAAGT	GCCTGTGTAT	TTATATCCTT
	TCAAATATCC	TTTGTAATAA	ATCATAAACA	TGTTTCCCTG	AGTTTTTATG	CCACTCTGTC
					TTTGAAGCCA	
	AGTTCTGGAG	GCCTGGACTT	GAGACTGGTG	TCTAAAGTGG	GCAGGCAGTC	TTGGGGACTA
5	AGCCCTCAAC	CTACGGGATC	TGAAACTGTC	TCCTGGTAGG	TAGCATTGGA	GTTGTACTGG
J	AGGGCACTGA	GCTGGTGTCT	GCTGCAGAAT	TGATTGCTTG	CTTGCTGGTG	GGGAGAACTT
	CCTACATATT	TTGGGGTCAC	CGAAGTCTTT	TGTATTGATT	GTTGTTGCTG	AGCTCCTATT
	CCTAGAAACT	TGCTTTATTA	AGTTCTGTTT	ATTCAGGGCT	TATCTGAAAG	AGAAACATTT
	тттатсатт	GAGATTTCTA	AGCCATTTTA	AAACTTGGTT	TTAACAGCTT	GAGAAATTGG
10	GGGTATAGTA	GGGATGGAGA	TACATATATT	TGGATGTGAC	TTCAAGCTAG	ATAAAAGTTT
10	GGAAGGAATA	AAAGTTTGAC	ATCTAAAGTT	TTTGCATAGT	TTGAGTTGAG	CAGGAGGTAC
	TAGGTATGTT	ТТТААААТАТ	TTTTTCAGC	CAGGCACGGT	GACTCATACC	TGTAATCCCA
					GCGTTCGAGA	
					TAGCTGGGTG	
15					ATTGCTTGAA	
13					CCTGGGTGTC	
	ААТААТААТА	AAAAAATGAA	AGATTTTTTC	TTACTCAGCA	TCCTCCAGGC	ATTTTATTAT
					ACTCAGGTGG	
					TCAGGCCTGT	
20					ACACTGGGCT	
20					GATTTTAGTA	
					TAAAAGTACC	
					CCTTTAATTC	
						AAATACAAAA
25	ATTAGCCAG	G CATAGCAGC	G CACGCCTGTA	A GTCCCAGTTA	CTTGGGAGGC	CGAGGCAGTA
23	GAATCACTT	G AACGCTGGA	G ACAGAGCAT	G GAGTGAGCTG	AGATTGCACC	ACTGCACTCC
	ATCCTGGGT	G ACAGCGTGA	G ACTCTGTCT	C CAAACAAAAC	AAAACACACA	CACACACAAT
	GTAACAACA	C GAAACAGAA'	T ACTGTGAAA	A TGCTTAATTA	TGTCTGACTT	TACATGATGG
						TTTTTAATGA
30	TGATCATGT	A TAAAGCTCC	T CTGTGTAGC.	A TTCTCTCCAC	CAAATTGCCC	AGAGACAGGA
30	AGTCCTGTA	а аасаааста	A GCTCCAAAA	A ATGACCTCC	GTTGAATAGG	CTTTTTTTT
	դարարարար 1101001011	T TTTTTTT	G AAATGGAGT	C TAGCTCTGT	TCCCAGGCCC	TCGCTCCTTC
	САССТССТБ	G GTTTAAGAG	A TTGTCCTTC	C TCAGCCTCC	A GAGTAGATTO	GATTACAGGT
	GCCCGCCAT	C ACGCCCAGC	T AATTTTTGT	A TTTTTAGTA	G AGATGGGGTT	TCACCATGTT
35						CTCCCCAAGT
33						A ACCTACTGTT
						A GCAGTAGAGA
						A TATGTAGAGG
	GGAAGTAGA	C TCATTATGO	C TGGAAATTI	T ATTTGGTGA	T AGTGAAGCA	A ATCTTAGGGC
40						A GCTAAAAAAA
						r TGGGAGGCTG
	GGGCTGGT	GG ATCACCTG	AG GTCAGAAGA	T GGAGTCCAG	C CTCAACATG	G TGAAACCCCT
						G TCTCAGCCAC
				A ACCCGGGAG		C AGTGAGCTGA
				22		

CATTGTGCCA CTGCATTTCA GCCTGGGTGA CAGAGTGAGA CTCTGTCTCA AAAACAAAAC
AAACAGAAAC AAGAACTCTA TGTTGAGAAA TCCATACTAG AGGGTTTAAT TTCTCACTTT
GTGTGCTAGT GATTAATAAA TTTCAAGCTT ATCACACAGC CAGAAATGTC GCCTTGTTT
CTACAATAAA AGATTTGGGA ATATTGTGAC ATTTTTTCAC TGAGCCTTCT GGGTTCATTA
TTAAGATATA ACAATTTTAG AAGACTTATT GAGATAGGTA TACTTTTTAA AATTTGGATT
CAATATATCA AGCTCAGCAT TTTGTCTTTT TTTTGTTTTTG TTTTAAACAA CTTTGGGTTT
ACTTGTAAGA GTATTTCAGT GGAGGAAAAG TATCAGAAGT TTAGCACTGA GTGCCTGCCT
AACTATTTA TGTCTTCTTT CATGATGATG CTTAATCTCA ACATAATGAC TCAGTTTACC
ATTGTAAACT CTTTGTAGGA TGTCACTGAT TTTATCAGTC ATTTGTTCAT GCAGCTGGGA
ACTGTGGGGA TATATGATTT GCCACATCTG AGGAACAAAC TGGGTGAGCT CCATTGAAAA
ATTTCTAACT TTCTTATTAA AAGCAAGCTT CCTCACTTCC TGATGGCTTA TGCAAATTAA
TGCCTATTTA TTCATCAGAG GGTGACTGGA GAGATTATGG TTTTCTTATG AATTTCTTGC
TTGAGCCTGC TTGTCTGCTG TTTCTAAGTT GTCTGCACTA TTTTATGTGA GTAATTTTCT
ACTTTATTAT GTTCTGTTTT CACATGTCAA ATCAGCTCTC CCAAGAATGC TACTTGTAAC
CTAAGTAGAC GTGAACCGAA AAGGGTAAAG ACCCAGCTAA AAAAAAGGTT GACTCAAGTT
CAGTTCACAT TCGTAAAGTA GTTAGGCTGC CTTAGGTTGC TTATTCTTTT CTTGAAAAGA
ATCCCTCAAG AGAGAAACAT GTGAGGCCAC AGCAGCTTAG ATCTGTCTTC CACAGAGAAG
GTGGCTTTTA CAGAGAAATT GACATACTCA TCACTTATCT GACATGACCC AGCTTTGTAA
AACTGGCTTC TATTAAAATA GTCTTAATAG ACTATTCACT GAGGAGGAGA GAATCTTATT
CACTCTTTAC ATTCTCTTCA CATTTTCAGA ATAGATGTTT AAATCATTGC TCACACTGGA
TCCATAAATG TCTAAAATGT TGATGAAGAA ATAGGTGATT AGAGAGTAAA ATTAATAGAG
ACTTACCCTT CCTTGCATTT TAACATAATA TTCTTTCCCC TTTTCCTTCC TCTGTTACTT
GGCTCTTAAA TACCAGAAGT GAGATATGAA AAAGGGAACT GGGAACAAGT ATTGAAAGCA
CCATAGGTTT ATCTTATATT AGCATTTTCC AAACTTTATA ATGAACCAGC AGTGACCCAC
ACATCTGCCA GGTAGGAATT GCTCATCAGT TGTGTCGCGT TATCTTGTTA AGCTTCAGCA
ATGTTCCATA CAGCCACTAA TAACAGATCA AAGTGAGCAT TAGGGTTGAA ATTAGTAAGC
TGTTTCTTCT CAGTTCTTTC TGGTAGTTGA ATAAATATAG AATGTATTAA ATAGTTTTCT
TTATTTCAGA ACTTCTCAGA GTCTGTAATA TATTGTATGG TGGTAGCTTA GGAGGAAGAT
GCAATAGGAA ACTTTTCCCA GATAGGTTCA CTATTTTTTT TTTCCACGAA AAATAAGCTG
TTCTCAAAAT ACAGTTTACA AAATTTTATC CTTAACTCTT CACTCTTTCT CCTAGTTAGG
GAGACCGCTC CACCAGTAGA AAAGATAAAC CCTGGTAATT TGTTGTGTAA ATGGGATAAA
TAGCCTAGTA CCTAGTCATG TGGATTCAGG CAGCACTGAG CCTAAATTAA AGTTTGCAAG
GTATACATGT TAATGTATCT AAGTTACTAT ATTTAGCCTG TTTCTTAAGT ATGTTTCAGA
AACATATTCG TTTTTTCAG TGGCAGTTAC CTTCAGATGC ATGTGCTTCT AAAGCATGTT
GGTTGCATGT GCAGTACATG TTGCTTAAGC ATTTAGCTTC AGAATGGCAT CTTTCCTGTG
AATGTCTTAA CATTTACAAA AATATACCAG GATCTCAAAT ATCAGTGCTG GTATTTTTTT
TTTTTTTTAC TTAAAGAAAC TGATATGATT AAATATTAAG AGACAATATG ATCCTTGTTG
GCTTGTAACC CTAGTTTTTA TTGTCTTGTA GTTATTAAAT AGAGCATCTG TTGAGGGACT
CTTTTAAAAC CACAGCCATG AACAGACGTT GGGGCTAAGA GACAGAGCAG CCTGCGACAG
TGTGGACCTA CCTGTAGCAG CTAGCAAAGG CCTCTAGCAG CTACAGTCCC TTCTGGAGTC
TTTATTTGCA TGCAAAATGC AAAGGAGTCC TGGTGACCTA CCTCCAAGGC AGCTGCCCTC
CTGAACACTC CCTTGGAAAA CAGTAAACAT CATTTTGGAA TGTGAACAAC CAGAGACTAC
ACAGGAGAAA GGAAAAAAA ATTCTGAAGA TGCAAAATCT TGGGTGGCTT CACCGTTCAG
TTTTTTAATA AAAGGAACAA TATACAACAC GTTGTTCTTT TTCTCTTTTG AAATCCCTTC

	·
	TATTACAGTG ATTTTTTCT AAGATTGTCA GGATTTGAAG TGTATGTTTT GTTTTATTCA
	CAGCGTAAAT TTTATTCACA GTTTAACTGT CTGCCTGAGT GTCTTTCCTT TCTCTAATTA
	CCTTGAGGAA CCCAAGAGCC TGTTGTAAGG AGAAAATAAG GCCCTTGGAT CTCTTGAGAT
	TCACAGATAT AAGTTATTGA AGGGAAGATG GTCCTATGGA GGACATATTT AAAGAAGGGA
5	AAAAAGAGGC TTTCTCAGAT ATGTCAGACT GCTATAGTAT ACTTCTACAG ATTATAGACC
	TCCAGTACCT CTGGCCAGAA AGATGGTATC GTAAACACCC TATTTTTTTT CTTTTTTT
	TTCATTAGGT ACAAGCTTTG TGCTAAGAAG TTGACATACT ATAAGCTACA AAAGTTCTGT
	AAAGTAGATA TAACTAGTTT CATTTTATAG ATAGAGAAAA TTAATCTCTT ACAGTGCTAA
	GCTCACAGAG TTTCTAACTG TAAAATGCTA GAACTTGTCT TTCAAGCCTA AAGACTTCCT
10	TGGGGCTAAA TAGTGAAAAA AGCCATTTCA CAAATAAGTA AATGGTATTT AGAGGCATAT
	TTGGATTTCC TGGTAAATTC CAGTCTGTGA GCATCATGAA TATTAGTTTA ATGTTGCATG
	GGCTCATGTT GAAGTTTTAA GAGAAGAACT GCCTTGAAGC TTAGGTTTCC TTAGCTATTA
	GGCTACTGAC TTTCTTGCCT AAACCAGGGT TTTTTCATTG AAGACCAAAA CTTACCTTCT
	CCTTCAGTTT GTAGTTTGGA AATTGGTAGA AGAGCTTTGT AAACTTCAAA TTAAGTACAA
15	ACTAAGTGTC ATAGTCAAAT TTACTAATCT TAATTACAGT ATTGTTCAAC TGATTGCTAT
	CTTCTAGCTC TTTCCTGCCG AATAATGGTC TTGTTTCCTG CTCTGTTGGT TTAGAGCTGA
	CTTCTTTCAG CTTTGGTAAG CCTGAAATTA TGGGGTTATG TTTAATTCAT ATTGTCTGGG
	TGGACTTTCC TCTCTTGCAT TTCTGCTTGA ATAGAAGAAT TTTTCTCTAG AGAGTAGTTT
	GTCATCCTTA CTCTGTTGAT TCAGATGACT CTTTGTATGA TCTGAGAGGT ATACTGTTCT
20	GCTATTCTGA GAAGAAGTAT TTCAGAAAGA TGAATTAAGA GTACAGTGGA CTGCTCCCAC
	CTGGAAACTT TTATCTATCT CACCTCTGGA CCTGATAAAT TCTTTATCAC TCAGGACCTT
	GATGACGCTG CTCTCTGAAA CCCTCCCCAG CTCTCTCTAT TACCGTGAGA AACATCAGAA
	CTTTGGTTCC CATTGCATAT CGCAGGTACC TCTGCTTTCA TGCCATGCTG TAATGGAGTG
	ATTGGGTAGC ATGTTTTCAT CTCTTTCCAG ATTGAAAATC TGTATTTCTC CCTGTATATC
25	TTCAACACCT AATGCACATA GAACTTTGTA GGTACCTGGA AAATGCACCA CAGTTTTCTT
	TTCTTTTTGC AGACTTTTCA CAAGTATTAC CAACTTACAA AGAATTAATT TTGTAGGATT
	CTAGAAAGAC AAATCAGGAA TGGTGCCATA TACATCTTTT TTGATTCCCT GCTCTAAAGA
	ATATTATCAG GTTACCTTCC TGCAGAGTTT TAAAAGAATT GCATATTTCA AGCTGACTTT
	CAGGATGTAA ATATAACCAA AGCAACTGAT ATGTAAAAAA TATATTCAAT GGCATTCCTA
30	GATTTTCTTC TAGGGTGTTT TATTGTTTTG GGTTTTACAT TTAAGTCTTT AATCCATCTT
	GAGTTAATTT TTGTATAGGT ATAAGAAAGG GGTCCAGTTT TAATTTTCTG CGTATGGCTA
	GCCAGTTCTC CCAGCACCAT TTATTAAATA GGGAATCCTT TCCCTATTGT TTGTTTTTGT
	ACGGTTTGTC AAAGATTAGA TGGTTGTAGA TGTGTGGTCT TATTTCTGAG ATCTTCATTC
	TCTTCCACTG GTCTATGTGT CTGTTTTTGT ACCATGCTTT TTTGGTTACT GTAGCCTTGT
35	AGTATAGTAT GAAAGATAGC ATGATGCCTC CAGGTTTGTT CTTTTTGCTT AGGATTGTCT
	TGGCTATACG AGCTTTTTTT TGGTTCTATA TGAATTTTAA AATAGTTTCT TCTAATTGTG
	TGAAGAATGT TAATGGTAGT TTAATGGGAA TAGCATTGAA TCTGTGAATT GCTTTGGGCA
	GTATGGCCAT TTTCATGATA TTGATTCTTC CTATCCATGA GCATGTAACG TTTTTCCCTT
	CGTTTGTGTC CTCTCATT TCCTTGAGTA GTGGTTTGTA GTTCTCCTTG AAGAGATCCT
40	TCACTTCTTC TGTATTCCTA GATATTTTAT TCTCTCTGTA GCTATTGGGA ATGGGAGTTC
	ATTCATGATT TTGCTCTCTG CTTGCCTTTT GTTGGTGTAT AGGGATCCTG GTGACTTCTG
	CACATTGATT TTGTATCCTG AGACTTTACC GAAGTTGCTT ATCAGCTTAA GAAGCTTTTG
	GGCTGAGATG ATGGGGTTTT CTAGATATAG GATCATGTTA TCTTCAAACA AAGACAATTT
	GACTTCCTCT CTTCCTATTT GAGTACGCTT TATTTCTTTC TCTTGCCTGA TTGCCCTGGC ${\cal A}{\cal H}$
	·

	CAGAACTCCC AATACTATAT TGAATAAGAA TGGTGAGAGA GGGCATCCTT GTCTTGTGCC
	AGTTTTCACG GGGAATGCTT CCAGCTTTTG CCCATTCAGT ATGATATTAT CTGTGGGTTT
	CTCATAAAA GCTCTTATTA TTTGAGATAC GTTCCTTCAA TACCTAGTTT ATTGAGAGTT
	TTTAACATGA AGCGATGTTG AATTGTATCG AAGGCCTTTT CTGTGTCTAT TGAGATAATC
_	ATGTGGTTTT TGTCTTTAGT TCTGTTTATG TGATGAATGA CGTTTATTGA TTTGCATATG
5	ATGTGGTTTT TGTCTTTAGT TCTGTTTATG TGATGATTGAC TGTGGTAGAT AAGCTTTTGG TTGAACCGGC CTTGCATCCT GGGGATGAAG CCAACTTGAC TGTGGTAGAT AAGCTTTTGG
	TTGAACCGGC CTTGCATCCT GGGGATGAAG CCARCTTONG TOTOTOGCTC GAAGTTCATC ATGTGCTGCT GGATTTGGTT TATCAGTATT TCATTGAGAT TTTTTTGCGTC GAAGTTCATC
	ATGTGCTGCT GGATTTGGTT TATCAGTATT TCATTGTGTATCT CTGCCAGGTT TTGGTATCAG AGGGATATTG GACTGAAGTT TTCTTTTTGT TGTCGTATCT CTGCCAGGTT TTGGTATCAG
	AGGGATATTG GACTGAAGTT TTCTTTTTGT TGTCGTATCT CTGGGTATCT CTGGGTATGGAA GATGATGCTG GCCTCATAAA ATGAGTTAGG GAGGAGTCCC TCCTTTTCAA TTGTTTGGAA
	GATGATGCTG GCCTCATAAA ATGAGTTAGG GAGGAGTCCC TGGTAGAATT CAACTGTAAA TAGTTTCAGA AGAAAGGGTA TCAGCTCCTC TTTGTACCTC TGGTAGAATT CAACTGTAAA
10	TAGTTTCAGA AGAAAGGGTA TCAGCTCCTC TITGTACCTC TGGTACTTCATA TCCATCTGGT CCTGGACTTT TTTTCATTAG TAGGCTATTT ATTACTGCCT CACTTTCATA
	TCCATCTGGT CCTGGACTTT TTTTCATTAG TAGGCTATTT ATTACTCTCT GGAGTGTGTA
	ACTTGTTATT GATCTATTCA GGGATCCAAC TTCTTCCTGA TTCAGTCTTG GGAGTGTGTA
	TGCATCCAGG AATTTATCCA TTTCTTCTAG ATTTTCTAGT TTCTTTGCAT AGAGGTGTTT
	GTAGTATTTG CTGTTGGTTG TTTGTACTTC TGTGAGATCA GTGGTGGTAT CCTGTTTATC
15	ATTTTTATT GTGTCTGTTT GATTCTTCTC TTATTTTTGA CAAAGCTGAC AAAAAGAAGC
	AATAGGGAAA GGACTCTCTA TTCAATTAAT CCTACTGTAT ATCTGGCTAG CCATATGCAG
	AAAATTGAAA CTGTTCCTGT TTCTTAATCC ATATACGAAA ATCAACTTAC GATGGATTAA
	AGACTTAAAT GTAAAACCCA AAATTATAAA ACCCTGGAAT AGAATATAGG CAATATCATT
	CTGGACATAG GAATGGGCAA AGATTTTATG AGAAAGACAC CAAAAGCAAT TACAACAAAA
20	GCAAAAATTG GCAAATGAGA TCTAATTAAA CTAAAGAGCT CTGCACAGCA AAAGAAACTA
	CTGTCAGAGT GAACAGGCAA CCAACAGAAT GGGAGAAAAT TTTTTCAATC TATCCATATG
	ACAAAGGTCT AACATCCAGA ATCTACAAGG AACTTAACAA ATTTACAAGA AAAAAGGAGC
	CCCATTAAAA AGTTGGCAAA GAACATGAAC AGACACTTCC CAGAAGATAT TCATGTGGCC
	AATAAACATG AAGAAAAGCT CAACATCACT GACCATTAGA GACGTGCATA TCAAAATCAC
25	AATGAGATAC CATCTCATGT CACAATGGTG ATTATTAAAA AGTCAAACAA CATGCTAGTG
	AGGTTGTAGA GAAATAAGAA CGCTTTTACA CTGTTGGTGG GAATGTCAAC TAATTCAACC
	ACTGTGGAAG ACAGTGTGGT GATTCCTCAA GGATTTAGAA CCAGAAATAT CATTACTGCA
	TATAGACCCA AAGGAATAGA AATCATTCTA TTACAAAGAT ACATGCACAT GTATGTTTAT
	TACAGCACTA TTCACAATAG CAAAGACATG GAATCAACCC AAATGCTCAT CAGTGATAGA
30	CTGGAAAAG AGAATGTGGA ACATAAACAC CATGGAATAC TATGCAGCAA TAAAAAGGAA
	TGAGATCCTG TCCTTTTCAG GGACATGGAT GGAGTTGGAA GCTGTTATCC TCAGCAAACT
	AATGCAGGAA CAGAAAACCA ACCACCACAT GTTCTCACTT ATAAGTGGGA GCTGAACAAT
	AGAACACATG GGCACAGGGA GGGGAATAAC ACACACTGGG GCCAGTCAGG GGGTGGGGGG
	TCAAGCTGAG GGAGAGCATT AGAAAAAATA GCTAATGCAT TCTGGGCTTA ACCCATTTAT
35	GCCTAGTGTT CCATTTCTGG AATGCTAAGC ATGTGGAAGT TCTTTATATC CTGCTCAAGG
	TCATTGCCAA GGTCTGATTT TTCACATTCA ACAAATTGCA ACCTCTGGCA TAAATGGGTT
,	AATACCTAGG TGATGAGTTG ATAGGTGCAG GAAACCACCA TGGCACATGT TTATCTATGT
	AAGAAACCTG CACATCCTAC ACATGTACCC TGGAACTTAA AAAATTTAAA ATATATATGT
	ATATATATT AATATGGAAT TTTAAAAATT ACTAATGAGT TCTTTTATCT GAGTAATTTT
40	GCATCAACAT GCTTTTATTA TGGAAGAGAA GATTCAGTGA GTACAAAATT GCAGATACAT
	GTGTCAGAAG ATCCCTGAAT ATAATAAGGC TTAGTATTCT GTGTCATAAT TGCCTGTTTG
	TATTCCTCTC TGGTCTTTAA ACTTCATTAG GGCAAGGATC AACTCCATCT TACTAACCAT
	TTGATTCCCT ATGTATTACA CGATATATGA CCAATAATAA GCCTTCAATA AATACTTGTA
	AAATAAAGAA TGTTATGTAA TATATCATGT GGTATTGTTT TATTGATGTG TTCTTTGAAG

	TGTTACCTTT GTGTCCTTAG AGGCTATGGT GGGCCCTCAG GCCATATATG TTGGATGATT
	TAGTGGAAAA GAGATTAGAC TTTTATATTT TATTCTATTT ATTTATTAT GTTTTAATCT
	TGTGTAAGCC TATAGAGGCC AAGATAGGGC TCAGAGTTAA AAACAGTGAT GTTCTAACTT
	TCAAAACTAC TAAAAACAAA ATTAGCTATC AAGGGAAATC CTGAGTGTTC TGACTGCAGG
_	GGTTCCCCAA AGCCTGGAGG ACCGAAGCTT TGAAATGCTG CATGGTGTGC CTCACCTCTG
5	TTGAGTATAG TGAGGAACAT GAGCTGAGGA GTGGATTAGA TGAACTCTTA AGATTTTTTT
	CCAACTTTAC GGTTTGAGTT TATGAGTAAG TAAGTGAGGA ATGAGCCCAG GGACTTTGAC
	TACTTTGCTG TGCAGTAAAT TGCTGTTTTC TTAGCTAATT CTGGAGTGTG TATTTTGGTT
	TACTTIGCTG TGCAGTAAAT TOCTOTTO
1.0	GTAACTTCCA GTTCCTTCAG CACTGGLZTT  GTAACTTCCA GTTCCTTCAG TAAAATGCTG GAAGTTGCTT GGCTGTAACC ACTGTAACTA
10	TTTCCTGGCA GTTTTGCATT TTATTACAAT TCGACTTTTT AAATGCATGG TTCAGTCCAG
	TTAATAGGCA GAAAACTATA GCTATTCCTT GGAACTGACA TAGGGAAGTT GGTTTCTTTC
	TTAATAGGCA GAAAACTATA GCTATTCCTT GGCAAAATAC CAATTCAGTA AAATTTGCCT
	TGAGGAAAGA TGTGACACTC TGATGGATTT GGGTATATA TAGATTTCTA CAGTAAACAA
	AGGGCAAAAA TGGAAACATT TTTTGTTTTT TAATGTTGCT GGAATATAAT ATTTCCATAT
15	AGGGCAAAAA TGGAAACATT TITTGTTTT MMTGTTGGT TTGGTTAATC ATCAAATCAT TCAACTCCAT GCAAAGTACT CTTTTCAAAA TTACAAATCA
	TTGGTTAATC ATCAAATCAT TCAACTCCAT GETTIGTTATTAT GTATTCTAAA TTATCTATAG TCTTGATGGA TTAGACTTGG ATGTCTAAAT TTTTTTATTAT GTATTCTAAA TTATCTATAG
	TCTTGATGGA TTAGACTTGG ATGTCTAAAT TTTTTTTTTT
	GTACATTTTG AAAATTATAT ATACATGTAT CACTOMOTH GITTER AAAATATATT ATATTAAGCG ATTAAATACT TTATTAATGT CATAAAATTG TGCCCTACCC AAAATATATT
	ATATTAAGCG ATTAAATACT TTATTAATGT CATAAAATTO TOOGGTTOO TTAAGAAGAT ATCATTTGAA AATAATGTAA AATATTCTTC ATTATTATGA TTTTGTTGCC
20	TTAAGAAGAT ATCATTTGAA AATAATGTAA AATATTCTTC MITMITTATTAAGAAGTAATT TTCAACATTT AAAAGTAATG TTAGAGTCCC TGTTACTGAT AATTTTCATG ACTGTAAATT TTCAACATTT
	AAAAGTAATG TTAGAGTCCC TGTTACIGAT AATTTICATG NCTGTAGAA TTTTGACACA GGAAACAAAA TTTTCTGATG TGTTTTAAAA TCTTAAAGAA TTTCAGTAGA TTTTGACACA
	GGAAACAAAA TTTTCTGATG TGTTTTAAAAA TCTTAAAAAA TTTTGTTAA AAGTATGAAA GTGTCACAGC TTTAGAAGGA TTTAGAATAA TTATCAATTG
	TTTTTGTTAA AAGTATGAAA GTGTCACAGC TTTAGAAGGA TTTAGATGT GTATTAGGGA TGACAATTTT AATAGACAAA TTTTAACTGA AACATTTGCT TTATTGATGT
	GTATTAGGGA TGACAATTTT AATAGACAAA TITTAACTGA MACATTTGG TTTTAAAAAAC CTAGGACTAA AAAAAATATT ATAGTGATGG AACCATTTCA AAACATCTGT TTTTAAAAAAC
25	CTAGGACTAA AAAAAATATT ATAGTGATGG AACCATTTCA TELEMONOO TOO TOO TOO TOO TOO TOO TOO TOO TOO
	TTCTGTCAAC ATGTTTGACA GAAGTCAGAA TAGGGTAATT MIZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZ
	TTATGTTTGC ATCATTAGTA TTTGCAAATC CTTGTTTCTT TGCTACTTTAA GTCAGTTTAA CTTGTTCATT TTATGAGTGT TAGCCTGATT AATTCAGATA AGATGCTCAA GTCAGTTTAA
	CTTGTTCATT TTATGAGTGT TAGCCTGATT AATTCAGATA AGMTGGTGATG GACTTACTAG GCAGGCAGAT GTGAAAAAGG GAATATGCTG TGCCTTAACT ATCTTATAGG GACTTACTAG
	GCAGGCAGAT GTGAAAAAGG GAATATGCIG IGCCITAACI ATGGGCACAC CAATGTTGTT
30	CACACTGTAC ATAACAGATG AACATCTGAT ATTAGATCTG ATGGGCACAC CAATGTTGTT
	TCATATATTA AATAATAGTG GAAGTTTTCA ATTTAATTAG TTTATTTATT TATTTATTTA
	TTTTTGAGCC GGTGTTTCCC TCTTTTTGCC AAGGCTGGGG TGCAATGGCA CATGGTTGCA
	GCTCACTGCA ATCTCCACCT CCCAGGTTCA AAGGATTCTT CTGCCTCATC CTCCCAAGTA
	GCTGGGATTA CAGGCGTGTG CCACTATGCC AGGCTAATTT TTGTATTATT AGTAGAGACA
35	AGGTTTCACC ATGTTGGTGA GGCTGCTCTC AAAATCCCGA CTTCAGGTGA TCCACCCGCC
	TCAGCCTCCT AAAGTGTTGG GATTACAGGC ATGAGCCACT GCGCCTGGCC AATTGTATTT
	TTATGTTAGA AATATATATG GGAGTTACAA TACTTTGTTT CCAATGTTCC GCTGGAAACC
	ACTGGTTGAA AGGATAGGTC CCAGTGTCAG CTGCCCAGTT GGCAGTTTTG GATAGAATAT
	AAGTTCTATG AAGGCATGGA TGAAGTATCC TTCATGGTAA TGATGGTAAT GTTAGTATATCTT
40	TCACTTGTTT AATGTTGTAT AACTTTTTAA CTCTTCCAAC TTCAATTTCA TCTATATCTT
	TCATAGCAGT CCAGTAAAGT AGTATCAATT ATAGATATCA TTCCTACTTT CAGATAAAGT
	AACTGAGGCT CACTAGTTTT GACTTGCTCA AGGTTACGTA AATGATAAGT GACAGACTGG
	ATTTAAAGGG CGGAGTCCAC TTTACTAAAC TGCCTCTGTC TTCTAGCCAA GTTTGTTATG
	TAAAGTAAGT ATTCAGACAT TGGCTAAGTA TATTATTAAT CAAGAAAACA GTAGAACAGG

10031E89 O1250E

PCT/US00/20400 WO 01/07663

	TTTAAGGAAT	TTTTTTTCC	CTCTCCTCAG	TGTCAAAAAG	CAGTCAAGTA .	AAAGATCCAA
				TCAATTCCAA		
				GGAAATTGAA		
				CAAATGGATA		
5				CATATTTTAG		
J				AAATTATCTC		
				AGAAAAATTT		
				TTGGTGATAC		
				CGAAAACATT		
10				GCGTACTTGG		
	TCTCTTTTAC	TAAGTTGTTG	TCGTGCAGAG	TGGCATTGCA	GCTCTGTACT	TTTACAGAAT
				TCAGATTGTA		
	TTTAAAAGGG	AAAATAATTC	CATCAATTTA	ATGGAATGTG	${\tt TAGGGTTTAA}$	GTTATAACAA
				СТСАТАТААА		
15				TGGCCGTGGT		
	GCACTTTGGG	AGGCCGAGGC	AGACGGATCA	CCTGAGGTCA	GGAGTTCAAG	ACCAGCCTGG
				ААТАСААААА		
	ATGCCTGTAA	TCCCAGCTAC	TCAGGCGGCT	GAGGCAGGAG	AATTGCTTGA	ACCCGGAAGA
	CGGAGGTTGC	AGTGAGCTGA	GATCGCGCCA	CTGCACTCCA	GCCTGGGTGA	CAGAACAAGA
20	CTCCATCTTC	AAAAAAAA 3	AAAATTAAGA	CAACCACAGA	TATAAGGATT	TGCAATTAGA
	AAGCTTGAAC	GGAAATTTTT	GGAACTTCTA	GGTCAACCTA	AAATTACAAA	TAGGGAAGCA
				GTAATACTGT		
	GTGATATAT	r GGCATCACCA	CACCTCTGTT	AATTATAAAC	TTACCTATTC	TCAGTGCCTT
				TTGTTCTTCC		
25				GCCAGATAGT		
				TGGAAAGAAC		
	AAAATAAAT	G AGTTAGTATA	CATAAAACAC	AGCCCATGAG	ATATAGTCTG	TAATTATCAT
						GATTCTGTCA
	TCCAGGCCG	G AGTTCAGTGO	G CTGGATCAT	r GCTCACTGTA	ACCTCAAAAC	ACCTGGCCTC
30	AAGTGATTC	C CCCAACCTC	G GCGTTCCAA	A GTGCTGGGAT	TAAAGGCATG	AGCCACTGTA
	CCTGGCCAT	T GCTCCTGTT	TAAAGATGA	G GAAACTGAAA	GTACAGAGAG	GAACGTGACT
						GCCTGAAGGC
						AAAGACTGAA
	AGAGCCAAG	G CAACATAGG	T ACCTGAAGA	A TGGGAGAAC	A CCTATGAGGA	CTGCTAGAAA
35						GAGTCAGTTA
						CTTTGTAACA
						A CCTTTTCTTC
						TAGGTGGAAG
	CCCTTTGGG	AAAAAATT TS	A GCCTTTTCT	G TCATTTCTG	G GTATCTCCC	TCAGTGATTT
40						A GAAGTGGAGG
						r TTTTTTTGTC
						A GTAATCTGTA
						A TAGACAATGT
	ATAAATGA	AT GGGCATGCT	G TTCTCCAGT	'а ааастатат 27	T TATAGAAAT	A GGCAGTCTTC
				~ <i> </i>		

	ጥሮር እ እጥጥጥጥ እ	GTTTGCTCAC	CCCTTTCCTA	TAGAACTCAG	GTTTATCATA	TACAATGTTA
		CTTTGAATAA				
		TGTTTTTATT				
		TTTGTGTATG				
5		ACTGCTTCCT				
5		TTACCAGCAT				
		CATAATACAA				
		CGATCTACCA				
		CACACAACAC				
1.0		CATCATATTG				
10		TGAAGCCATT				
		GGGTACAGTG				
		CTCAGGTCAG				
		ATACAAAAAT				
4		AGGCAGGAGA				
15		TGTACTCCAG				
		ATCACCAAAG				
	TATCATGAAT	ATCACCAAAG ATATAACACA	ATAAAAGCAA	ATTCCCACAA	TTCAAGTCCA	GTTTTTCTGG
		CATACATTT				
	ATTCTAAAGC	CATACATTTA GTAATTTTAA	CERTAL ACTION	7.7.7.4.4.4.4.4.4.4.4.4.4.4.4.4.4.4.4.4	ACACATTTCT	CTAAACTGGG
20		GTAATTTAA CTACAAACCT				
		CTACAAACCT GTATTCCCTT				
		CTAAAATATT A AATAATGTGT				
25		A CCCTACTGTC				
		A TTTGTTTGAG				
	AACAGTTGC	A AGAGAACTAA	GIGGCCIGCE	AAGCCIAGIG	CCCACAATCT	
		G TTTACCAAAC				
		r ttttctacti				
30	CTTGCCCAA	G GTTCTACAA	TAGTCTTTTT	TATTAAATTA	AGICAGAGCC	CTCCCACATG
	TGTGCCCAA	G TCTCTCCTC	A GAAATTTAGO	C ACAGCATTTT	CCCTTCIGI	A A CATCAGG
	CTACACAGT	C ATCCTGGAT	ATGTGCCCCC	G CCTCCATGAC	GAGAIGIICI	AACATGAGGG
	TGACTAGGC	T CTGGAGCCA	G AGCATCTTAG	TTCAAATCC	C GGATCTGTTA	AAAAIAAIAG
	TTATTTTTC	A GTTCTAACA	TGAAATAAG.	A CAAAAGGCCA	TATTACTGAA	MCCTN ATA A A
35	AAGAATATA	T TTAATACTC	A GGTTACTTT.	A GCCCCCTCA	COMONGACIO	TOCIAAIAAA
	TTGAGAGAT	T GACAGAAAA	A AAGGATTCT	G TTTCACTGT	GCTCACACC	CCCTCCCCAA
		A GGCCGAGGT				
	CATGGCAAA	A CCCCGTCTC	т астааасаа	T ACAAAAATT	A GCCAGGCGTC	B MCCCACCTCC
		C CAGCTACTC				
40						AGCAAGACTC
		AAAAAAAAG				
	TCAGTGGA	AA ATTGTAGTG	T GATAGACTG	A GGAGAATTC	T GTTTCTGCC	A CTTCGCCTGT
						G TAAAATGTGG
	TTATTATC	AT CTGCATCCT	C ATAGTCTTC	T TGTAAGGAC 2 <b>9</b>	T GTGAAGTCA	C TTTTGAAAAA

	TGCCTTGAAT	ATGCCAGATG	GTCTGTTTTC	TTTCTTTTTT	GTTTTGTTTT	GTTTTGTTTT
					TTTGTTTGGT	
					TAATTCTTAC	
					GTGAATGAGA	
5					TTCTCTTATT	
J					GACAATGACT	
					GAAGGTGGGA	
					GGAAAAGTGA	
					TATGTGTGTA	
10	ATTTTAGTTT	GATTATACAA	AATATCTTTG	GCTTAGAAAT	ATTAGGCATG	CTATGTAAAA
	CCTTACTGGA	AAAATAAATT	GACCAACATT	ATTGAGAGTA	TTTTTTCAAA	GTGTTCCAAA
	AGTAATGGAC	CAATGATTAC	TTTAAATGAA	ATCATGTAAT	GGACCACAGA	ATTGCAAATT
	ACTAATAAAG	AAAAGCCATT	TTGCTTATTG	CCATGTAATA	ACATGTTGCA	TGATACAAGT
	AGATACGTAT	GTTTATGCTG	CAACAAGTAT	AGGTGATACT	AATTGGGCAA	CTTTTAAACA
15	AGACCATAAA	TAACTGAAAT	CAAAGTTCTT	AGTATTTATG	CAGCCTGTTG	GTTTGCGAGG
	GCTGCCATAA	CAAAGTACCA	CAGACTGGGT	GGCTTAAGGA	AGGTTGTTTT	TTCACAGTTT
	TGGAGGCTAG	AAGTCCAAGA	TTGAGGTGTT	GTCAGGTTTG	GTTTCTTCCA	AGGCCACACT
	CCTTGGCTTG	CAGATGTCTG	CTTGCTTACT	GGGCTCCTAA	ACAGTCTTTT	CTGATTGTCC
	TAATGTCCTC	TTCTTATAAG	GGCATCAGTC	ATATTGGATT	GGGGCCCACC	CATATGACCT
20	AATTTTACCI	TAATGACCTC	TTTAAAGCTC	TGTCTCCAAA	TACAGTCATA	TGGCTGTGAT
	ACAGGGGGTT	r AGAATTTCAA	CGTATGAATT	TTGAGGGGGA	CACAATTCAG	ACTATAAACT
	GCAGTTAATO	TTTACTGTTA	AATTAAATCT	' ATAACTGACT	AAACCACACA	AACAGGAGAT
	TTCAAATTAG	ACTTTATTAC	TTTTGGAAGG	AAGAGATGAA	ATGTTGTTAC	TTTTTCTGTT
	TTATGTAGT	r gtaaaagcca	CTAAAAATGT	ACATGTACAA	ATCATCCCAA	GCCAGGCAAC
25	ATTAGTAAT	G AATGACTGCA	GCACAGAGGT	AAGAGAGATG	ATTTAAAGGA	GGATGAATTG
					CTTCATTTTA	
					TACTTTTAAC	
					CTATTTGATA	
						GCCTCACCTG
30	TGAAGTGAG	C ATGTGGGAC	r GAATTGCCT	r TGAGGTCTCT	TTCAGCTCTT	AATGATCTGT
						AAAAATAATA
						ACACACGAGG
						CAAATATCTA
						TTATTTACAA
35						TTTTTGTTAT
						GGGTACGTCT
•						TTAAAACATT
						TAGACCATAT
						A GTATGCCTAA
40						GAAGTCTTAG
						G GTGAGTTCTA
						A ATAATCAGCT
						r TACATATATA
	TCCAAGCA'	TT TGGCACAAC	G TAAACCAAT	G TGGCTCAGA	T TTTAATGTT	A TTGTGAATCA
				- /		

	CTTGGGGATC TTGTTTAAAA AAAAAAAAA AAAGGAGATT CTGATTCAGG TGATCTGGGC
	TGAGTACTTA CAGTATGCAT TTCTAATAAA CTTCTGGGTG TTGCTACTGA TATTGGTTCA
	AGAACTAAAT TCTGAGTAAC AAGTATAAAC CGAATGTCCA TAAAGTATAA TTTCTTTTAG
	TATCATATGG TTATAATGCT TGCATTTTGT ATATATGTTG CATACAGGTT TTGCTTTTCT
5	CTTTTTTCTT ATTAAAAAGG TTATAAAGGT ACACTGTAGA AACTGTAGGG AGTAGGGGAA
J	AGCTTGGCAT CTCATGCCAC CATTCAGAGG TAACCATGTT TGGCATTTTA AAAATGCATT
	TATAGTGTGT GTGTGTGTGT GTGTGTGTGC GCGCATACAT ATATATGTAT
	GTATTATACA TATACGTTAA TATACATATA ATATACATAT ATTACTGTGC AAAAGTTGAT
	AAATTGATTA TGATTATGTA TAAACTTTGA TATCTTCTCA TATTTAATAT AATATAGTTG
10	TGAGCACTCT TCCATATCAT TAAATAACTT TCTAGACATT TTAAATGAAT ATATAGAAAA
10	TGAAACATCA TTAATCAGTC CCTTATCTTA GAATATTTAT TTCGGTTATT TACAATCTAT
	TAGTAGTTTA AAAGTACTTC ATGGAAAATC CTTATACATG CATCTTAGTT AACTTGAACT
	TTTATTTTTA GAGAAGACAT GCCAGGCTGT TCTAGGCTAT TGATACATAT TGCCAATTAC
	CTTCCTGAAA GATTGTACCA CTTTCCACTT AGGCACAGAG TGTGAATGCC TGTTTTAATT
15	ACTCTATTTA CATCAGTAAA GACTATATCT GAAATTATCT CATTTTTGGG GACAGGTATT
	CTAATAGGCT CTACAAGTAT GATCACAACT TTGTGAAAGC TATCAATGCC ATTCAGAAGT
	CTTGGACTGC AACTACATAC ATGGAATATG AGACTCTTAC CCTGGGAGAT ATGATTAGGA
	GAAGTGGTGG CCACAGTCGA AAAATCCCAA GGTAATCAAG CACACATTTT ATCATTAATA
	AAAATATGAA TGCTGAATAC CATCCTCCCT CCTAAGCAGT CCTACCAGTG TTGCTCGCCA
20	TTTTATTGGT CATCCCAGTT GTAACTTTTT ATGTGATCTG TTAACGATCT TTATCTCCTA
	TTGATACATA TTCTGTCACT CTCCAAGTCC TATTTATGAG TTTTTCTTTC ACATGTCTGT
	TGTTTCCATT GCCCCCTTTC CATTTTCTAC ACTCTGGCAT GCAGCTTTGT CCTGTTTCCC
	CCATGCTATG GATTAGCCTC CTGCCGGTAA CCACCTGAGG GGTTCTTCCT GCCTGCA
	TAAAGAAAAA CCATGGCAAT ATAGTAGAGA AAGAGCTTAA GAGACACTAG GCTGGCCATG
25	CCACGTGGGG GATGGAGTTC ATATTCAAAT TATCTTGTCC AAAGCTCATA GGTAGGGGTT
	TTTCAAAGGC AGCTTGGGGG AAGGGGTGGG GGTGGCCAGG TAACAGATGC TTGCTGCTGA
	TTGGTTGGGG TGGGTGAAAT CACAGGGAGT TGAAGCTGTC CTCCTGTGGG CTGAATTGCT
	TCTAGGTGGG GCCATAGGAG TGGGGTTGCT GGGTCCAGGT AGAACCACGG GTGTCAGACA
	TGCAAAAATA AAATAAAATA AAATAAGATA ANGGTAAGAT AAAATAAAAT
30	AATAAATAAA ATAAATATAA AATTTCCCTG AAAAGATATT TCAAAAAGCC AGTCTTAGAT
	TCTACAATAA TGATGTTATT TGCTGGAGTA ATTGATGGAG TTGCATGTCT TATAACCTCT
	GGAATAACGG CTGACAATCT CTCAAGTCTG CGCCTTAGCT GGACTCAGGT TCCTCTTCTC
	CCCACAGCCT GACTGCCTCC ATTAGCTTCA CAAAAGTGGT TGGGTTTCAG GGCAAGGCCC
	ATTGTCATTT AAACTGTAGC CGAAATGACT TCCAAAGTTA GCTTGGCCCA ATAGCCCAGG
35	AATATTTAAG TGGAAGGCAA GATGGGGGAT GGGTTAGCTT AGCTCTCTTT CACTCTCATA
	GTTTTCTCAC TGGTATAATT TTTGCAAAGG CGGTTTCATG CCTGCCATCT CTTTCGTCGC
•	TACCTCTCC AGTTCCCATT CTTAGCTGTT TTATGAAATG CTTCTAGTTT CATCCTCTTA
	TACCAAGTTC TGGGAGACTG ATTTGAGTAA TAATAAAACT CCAGTTTCCC ATACAGCCGG
•	CTCTGCGTGA ATTAAACTCA TTTTCTATTG CAATTTCCCT GTCTTGATAA TCAGTTCTGT
40	GTAGGCCGTG AGGAAGGAGA ACCCGTTGGG TGATTACGAG ACTGTGTTAC TGCCCACTAC
	CTAATGGATA CATTTAGCCT GGTATCCAAA CCCATCTAAT TATGACCATA ACTATATTA
	TCACCTTGCT CTGCTTTCAA ACGATATGAC ACACAATGAA TGAAAACTTT CATTTTCAT
	CTTCATTTGT GCTGTTCCCT TTGCCTCAAA TAGCCCTCAA CTTGCCTACA GTAACTGTAA
	AATTTGCCAC CTAAAAAAA ATCTCAAAAT CCTCTCTATG CTTTGATGTC CAGCAAAAAA

PCT/US00/20400

### WO 01/07663

•		
	AAAAAAAAA AAAAAAAGT TATCAGTATA GCTCTATTAC T	
	TTTCATCTTT ACCCATAAGC CCCATGGTAT ATTTCTTAGT T	TTGGGAAATT ATATCATAAT
	ACTTTGAATC TGTCTAGTCA GAAATTTAGA TAATTTAAAA A	AAAATTATT TTTCTGAAAG
	TATAGCTGGA TCTAGCTTAG GATCTACATC TATCTAATAT A	AGTTCCTTAA CATNITGTTA
5	AATGGCCACN GGNATAAGTC CTGTAATGCC ATACTTTGCT	TTAGGATCAT GTGACTAAGG
	GGTAAGGAAT TGGAAAGCAA TGGGGAGCTA GCAGAATTTG	AAAAATTATA TAGGTAGGTT
	ATTTTTCCTT AATACATTGA AATAGCCTCA AATTCTCAGA	GAATACAATG TTTAATCCTC
	TAATATCTGT AGAGTTCATG GCTCAAATTT GTATTATTTG	
	ATAATTACTC AACTAAAACA CTTTGAAATG TGAGAGTTCC	CTTATCTCCC TCGCAGGCAT
10	ATGACAGAGG TGTGGCTTCT CACACCTTTG GTTGCCCTTG	
	GGGGGAGCAT GCAGAGGGGC AGGTGCAGAG GCCATGGGGG	
	CACAGCAGTG TCTAGGAGTG GATGTTGGAG ACTCCTGAAG	
	AGTGTGCTCT TTCAGCTTAG CCGTCTGCAG ATGGCTTGTG	TTAATCAGGT CATTAGACCC
	CATGCCTTAT TGCAAGGGCA GGGGTCCAAT GTGACAGCCT	AAGTTCTTGC TCAGTGTACC
15	AGAAGAATTG GATCACACGT GGGCTGGAAG GATGAGCACA	AGGTTTTATT GAGTGGTGGA
	GGTGGCTCTC CGCGAGACGA CTCTCAGCCA GAGAAGGAGG	
	TCCCTGGAG TCCACTTGTT CCTGTTCTCC TCTGGGTTCA	AATACCTCTT CCCTCCTTTT
	CTGCTGTGCT GTCCCACCAC TCTCCACAGC TCTGTGCCGC	TCTGTTCCTC TGCTCCTCTG
	GATGTTCAGC TCCTTGTATC TGTGCCTGCT AAGATTTTGG	GTTTATATGG GGGAAGGATG
20	GGGGGCATGG CGGGCCAAAA GGCACCTTTT TTGGTGTGAA	AACAGAAATG CCTGTCTTCT
_ •	CTTAGGGCCT TAGGTCTTCA GGCTTGAGGG TGGGGCCTTT	
	TACCCAGTAT TTCCCTGTCT CCTGTCCATA TCAACATTAC	TAACTTTTTC ATCTGCAGAC
	TAATAATGCT AAGGTGTGGC ATTTTTCAAC TGTGAGACTA	TGTGAAGGTT TTTCCTGTCC
	AACTGATGGC ATCCTCCCAT AATTCTACCC CTTTCTTAAA	AGAATCTTTT GCAGTATTTC
25	TCCAAGTTTA TTCTAGAGAA TTTCTTGTCT GTGAAATGCT	CTAGTTAATC AAAATTGGAA
23	AACGGAGCAT ATCATATCCC CTTCTCAAAT TCACCAAAGT	GAAGTCCTAA TGTGTCTTAA
	TGTATCTGCA TGAGACAGGA AGCTGAGATC TATTCAACAA	CAAAAATCCA AACAAGCATC
	AAGAGGAGGA GTGTTAGCAC TTGAGCCTAG GGAGACTGTG	GCTCCTGCCT GAAAGATGGG
	AGCCTCAGTC ACAGCTGCTT TACCAAGTGT CATATGCTAT	GTTTCTGAGG ACTCCTGCTA
30	AAGCTCCCTT CTCCCTCCAG CCAACCACTT TTGTTTTAGA	CAAGGGCTGG GTTTATGAAG
50	GACTGTTTTC ATGACTAAAG CTTTATAGAA GGTTTAAGAT	AAGGAGATGG AATTGAGTGA
	AGTAGGAAAT ATGAAAGCAG ATATTATAAT CTGGCTTCCT	GATTTTTCAC TAGCATTTTT
	GTTTATAAAT TAGTTCTGTT CTAAGAATCC AATGACGTAA	TAGAAACTCT CAAAGATTCT
	TAACTTGAGA TATAGGGAGT CTTTGAAACT GCTGAAATTA	CAGACAGCAT TTATTGTTTA
35	TGAGCATTTC TGAATCTAGA GCTTTCACTA GATTTGTAAA	
55	AGAGCCAATC CTGTATCTTG TACTCAAAAT GTTTGTAATT	CTCACCTTTT TATCCCCCAG
	ACTCCTCTTG CTCTCTTC ATTTTCACAG TGTTACTGGA	AAGGGGTCCT GATCCAGACC
	CCAAGAGAG GCTCTTGGAT CTTGTGCAAG AAAGAATTTA	GGGCGAGTCC TCAATGCAAA
	GTGAAAGCAA GTTTATTAAG AAAGTAAAGG AATAAAAGAA	
40	CAGCCCTGAG GGCAGCTGGT TGTCCATTTT TATGGTTATT	TCTTGATTGT ATGCTAAACA
<b>4</b> U	AGGGATGGAT TATTCATGCC TCCCCTTTTT AGACCATAGA	
	CATGCATTC TTTTTTTTT TTGACAGAGT CTCACTGTGT	
	AAAGCAATCT TGGCTCACTG CAACCTCTGC CTCCTGGGTT	
	GCCTCCCAAG TAGCTGGGAT TACAGGCATG ACCCACCATG	
	3/	

	TTAGTTGAAA CAGGGTTTTG CCATGTTGGC CAGGCTGGTC TTGAACTCCT GACTTCAGGT	
	TTAGTTGAAA CAGGGIIIIG CCAIGITGGG GIGGTGGGGGGGGGGGGGGGGGGGGGGGGGG	
	GATCTGCCTA CCTTGGCCTC CCARROTGCT GCTGGTGGGA GTGTAGCAGT GAGGAAGACC	
	AGAGGTCACT CTCATCGCCA TCTTGGTTTT GGTGGGTTTT AGCTGGCTTC TTTACTGCAG	
_	AGAGGTCACT CTCATCGCCA ICIIGGIII GGIGGGTCTTTATG ACCTGAATCT TGTGCTGACC TCCTATCTCA	
5	TTCTATGACT TAGAATGCCT TAACTGTCTG GGAATGCAGC CCAGTAGGTC TCAGCCTCAT	ı
	TTCTATGACT TAGAATGCCT TAACTGTCTG GGTMTGCTGG GGTMTGCTGG GCTATCAA GATGGAGTTG CTCTGGGTCA AACACCTTTG ACAACATCAT	•
	TAAGCCTCAG TTCTCACACT GTTTTTGTTT TGTTTGTGTG TGCAATGATG GGCAAATCTC	:
	TAAGCCTCAG TTCTCACACT GITTIGITT TGTTTGTGTG TGGCCAGAT TGCCTTATAG GATGGTAGAA AAAAGGAACT TAATATTGTA ATGACTGTTT TGTGCCAGAT	•
	AATTGCTTTA AACTGTATCA GCATCTTATT TAGTCCTGTT AATGATATGA ATGTTATCTT	
10	AATTGCTTTA AACTGTATCA GCATCTTATT TAGGGCTGTT TATTGCTGTTCT AAGCACTTGT	
	CATAACAGCT GCCATTTAT TAAGGACTTA TCAGAGATTT TOTAGAGCATGA TTATACCCAC TACCCATTAT TACATTGAAT TTTCATAACA ACCCTTTGAG GTAAGCATGA TTATACCCAC	:
	TACCCATTAT TACATIGAAT TITCATAACA ACCCTITIGAG GITAGGITCO TITCACACAGAT TTTAATAGAA GTGAACTGTA GTTTGTAAT GTTAGGTTCC TTGCCAAATG TTACACAGAT	r
	AGTAAGTGAT AAAATCATAT GCCCTGAAAT TACATTATGC TGCCAAACTT AAATTTCTTT	r
	AGTAAGTGAT AAAATCATAT GCCCIGAAAT TACATTATGC TGGGTZTGTTA ACACATACTA TTTATCCTTT ATATTAGTAT ATTCTTAGGT TTAAACAAGA CAACTAGTTA ACACATACTA	A
15	TTTATCCTTT ATATTAGTAT ATTCTTAGGI ITAAACAAGA CIICINGTIO GATTTTGTCC ACAGTTCCTG GCTCATAACT CCCATAGCCC TTGTCACTAT CTTTTAANCC	3
	GATTTTGTCC ACAGTTCCTG GCTCATAACT CCCATAGCCC TTGTGTGTGTTCTCTCTCTCTCTCTCTCTCTCTCT	2
	TTGGGGCACA TTAGGCCTCA GAAGTAGGCC TCANGAAAAC AGAGTCTCTCA TCTTCTCCTC	3
	TCTCTGATCT TCTTCCGCCC TCCTCTCACC TGCCCAGGGC AGCACTTTAA TCTTCTCCTC	A
	CCTTTCTGAT CTTGGGTCAT AAGACCTTCA TTTCCAAAGA TGTCCTGTGT CATACCCTA	A.
20	AGGAAGGAAC ACTGAACAGA GAGAGGCTCA GAAGAATCTG GACAGGCCTT GCTGTGTTT	G
	CATCATTCCC TTTATGTCCA GTCACATCTC TACATGGTTG TCAGTTGTGC CTATTTGATG	G
	AAGTCCCCAT ATAAGGCTCA CAAGGACAGG GTGCAGAGAG CTTCCAGATA GCTGAACAA	C
	TGGAAGTTCC TGGAGGGTGG CGTGTTCAGG GAGGGCATGG AAGCTGTGTG CCCCTTCCC	c
	CATACCTTGC CCTACTCATT TCTTCATCTG TTTCATTTGT AGTATCTTTT ATAATAAAC	.C
25	ACTAAACATT AGTTAGTATT TCTCTGAGTT CTGTGAGTCA CTCTAGCAAA TTAATTGAA	A
	CCAAGGAGGG TGTCATAGGA TCCCCNACAT TATAGCTGGT TGGCCAGAAG CACAGGTAA	'C
	CAACCTAGGG CTTTCAATTG GCATGAGAAG TAGGGGGCAG TTTTGTGGGA CGGAGCCCT	'C
	AGCCTGTGAG ATCTGATGCC ATCTCTAAGT ACACAGTGTC AAAACTGGAT TGGAGGACA	'G
	CCAGCTAGTA TTCACTGTGA AATTGGTTGC TTGCTTGATT TGTGGGGAAA AACCCACAT	λA
30	CATTTGATCA CAGAAGTCTT TTGTGTTGAC AGTTGATAGT GTTCAGTGAG AGAATTAAA	'A
	AAAAATTGAG TTTCTTCTTC AACATACTCT CTCAATGTGA AACCACAGAA ACTATTTCC	ΔT
	TTCAAAGATG GAAATGGTTT GTTTGCATCT TAGTTTTTAT TTATACATCT TAGAAGAAA	יידי
	GTCCAAGCTT TGTTTTTCT CTCACCCTAT ATATAAAATT ACCTATGAGG CACAGATTT	ىلەر -
	TATGATCCTT GATTATAG ACTTTGTCCA AATTGTGTGT TTTATAGCAT TACTGTAAC	22
35	TGTTATAGTA ATCTTTGTGT ATATTATGTC TCTTAACATT GTCTTCCATA TTGTTAATG	מב
	CCATCTCATA TTTATCTCTG TATCATGTAT ATCTTCAACC AATGTGACTG GCTTAGAA	<u>አጥ</u>
	AAAAATTAGT GAACAATTAA CTAGTGTTTG TGTAATCTAT ACAATTGTCA AGGTTACAA	<u>م</u> لا 1
	TGCTATTTTT GAAGAAATCG TTGTTGTTTT TCTCTTTGTT TCATCTCAGT TCCATTTC	CC
	CAAGGATTCC TTTTTTTTT TTTTTTTTT TTTTTTGAGGC GGAGTCTTGC TCTGTCACG	GC
40	GGGCTGGAGT GCAGTGGTGC AATCTCGGCT CGCTGCAAGC TCCACCTCCT GGGTTCATC	ag
	CATTCTCCTG CCTCAGCCTC CCGAGTAGCT GGGACTACAG GCACCCGCCA CCACGCCC	CT
	CTAATTTTT TTTGTATTTT TAGTAGGGAC GGGGTTTCAC TGTGTTAGCC AGGATGGTC	TG
	CAATCTCCTG ACCTCGTGAT CCGCCCGCCT CGGCCTCCCA ATGCTGGGAT TACAGGCG	GТ
	AGCCACCGCG CGTGGCCCCT TGTCATGTAT TCTTAACCTG TGTTATATCT AAGAGAAG $3$ 2	

	TOTAL TOTAL CARACTECTAT GAAGTAGCCC AGAAAGAATC
	GTGAGAGGCG GGACCATTTG TGGATAGGTG CAGAGGGCAT GAAGTAGCCC AGAAAGAATC
	TTTGCCATTG ACTAAATTTT AGCCTAGTAA AAAACATGTG GTAACCACTG AAAAACTAGA
	ACAGTGTGAT ATGACAATGC CCAGTTGAAT AAGAAATTCA AATAGTTTAT AGTAACAAAA
	AATAATTTT ACCACATTGT GACTAGTGCC CTAAAAACAA ATTTGAAGGA GCAGAGAGAG
5	AGAGGAAGTG AGTTTTCGCT GGGGGCTGGT GAGTGGAGGC TACTTTATGA GCAGTTTTGA
	AAATTACAAG TTAGGGAAAA ACTCTTAGGG GAGTTTTAAG TGGTGAGCAA CAGGCACTTG
	AGGTTACAGG AGCAGCAGCA TAAAGATATA GAACAGAGAG GCCATTTGGC AGTCTTGGGG
	AAACTCCAAG AAGTCGAATG TGTCTAAAGC TGTGGTTCTC TCTCTGTGTG TGTGTGTGTG
	TGTGTGTGTG CGCGCGTGTG TTTGTGTGTG TGTGCATGTG TATGTATGTG TGAAATCGCT
10	AATACAGAGA AGCTGTTAGA ACAATTTTCT CACTTCCCAA AAATGTCCGT TCATTCATGG
	GCCTAGGCAA CTCTCCTTTG TGTGTCTTAC AGCCCAACAC TGTCATATAA GGTGTGGTTC
	TATAATGAAA CAGTTTTATC TGTTGTTTAC GGAGCTAGTC AGCCTGTGTT ATGCTTGTTA
	CTGGTTGAGG GTGTCCAGGT TCTTGGCTTC TTGAACAAAG AATTGGACAA AACTCACAAA
	TGAGGCAAGG AAAGAATGAA GCAACAAAAG CGGAGATTTA TTGAAAATGA AAGCACACTC
15	CACAGGATGG GAGTGGGCCT AAGTAAATGA TTCAAGGCCC TGGTTACAGA ATTTTCTGGG
	GTTTCAATAC CGTCTAGAGG TTTCCCATTG GTTACCTGGT GTATGCCTTA TGTAAATGAA
	GAGGGTGGAG TTAAGTTACA AAGTCATTTA CTCAGTATAG GCCTTGTGTT AATGGAGAGG
	GTGTTACTCC TGGGGGTTGT GGCCCATGTA AACGGAGAGG ATGAAGTGAA GTGACAAAGC
	CCTTCGCATT CCTGCCATTG CTGAAGTGTT TCCACTTTAT TTAGTTCTAG GAAGTCAGTG
20	TGAATTGGCC TTATGTTCCC TGCCTCCAGA ACCTGTTCTC CTGCCTCACA TAGACCCTTT
20	TTTCCCTTTG CCAACTTCAC TTCTTTTACA GCCACCTCAC TACCAATGTG TCTGTCTCCT
	AAGTCAAACA TCAGTGATTC CTCTGTTTTC TCTAAACCCT TCTTATGTCT TCTACTTCTC
	ATCTCTTTCT TGGTCAAAAA TCTTTCAAAA ATGAGTAAGA ATGCAGCTAT TCAGGCAAAC
	TAAAAATAAC ATCACAGTGA TATACAAAAC CAGTGTCATT TCACAAAGGA AAATTATCAA
٥٠	TAAAAATAAC ATCACAGTGT TAAAAAATAAC ATCACAACTCC ATTGTGTAAT TACTAGATCC TGAAAAAGAA ACAGCGAATG AAAGCCATTT ACACAACTCC ATTGTGTAAT
25	TACTAGATCC TGAAAAACCT TOTAGAT CTCTGGGTCT GAATCTAGAT CCTAGCTAGT TGACACATTG AATCACTCAT AAAACAGGTG CTCTGGGTCT GAATCTAGAT CCTAGCTAGT
	TGACACATTG AATCACTCAT MILLIONGOOD  CTGGTAGCTG AAATCATAGA ATTATAGTAG AGTTTAGGAA ATCATCCTCA AAGGAAAGAT
	TATATGTTGA TATCAAATGT ATATTTCCTT TCTAGGCCCA AACCTGCACC ACTGACTGCT
	TATATGTTGA TATCAAATGT ATATTTCCTT TOTAGGGGGAAAA TGTTCATGGT GAAATACAGC AAAAGATTTT GCATTTGCCA ACATCTTGGG ACTGGAGAAA TGTTCATGGT
	GAAATACAGC AAAAGATTTT GCATTIGCCA ACATCTTOOS TOO TOO TOO TOO TOO TOO TOO TOO TO
30	ATCAATTTG TCAGTCCTGT TCGAAACCAA GGTAAAAAAT TTTGGGGGGATACCTAG ATTTGTTTGG AACTATTTAT TGAACAGTTG CTCTGTGTGA TGGATTTCGG GGATACCTAG
	ATTTGTTTGG AACTATTTAT TGAACAGTTG CTCTGTGTGA TGCATTATTC
	ATGGAATGGG CATGATCCCC TTTTTACAGA AATAGAAAAT AGGTGGCCTA TGAATTATTC
	TTTCCTTTTA TATCCATGAC AAACTTTAGT AAAAAATTTT CTTTTCTACT GAGCTTAGCA
	TTTATTCAGT ACTCTTCTCA ATATATTTTC CAGGTAGCTA GTGACAATCA GAGTGATATG
35	TAAGACAAAC TCATTTGTCC TCCTAGTAGG AAAGATTCTA GTAGAAGCAA AGAATTGTGT
	ACCATTCTGC AAGTGGTTTG TTGGAATCTT TCTTTGATAC CTGTTCCTGT ATTCCCCCAC
r	CCCCATTAAT TTAGTCATTA ATTACTACAT GAACCCGTAA AATAAATCCT TAAATTATTT
	TCCTGGTAGA TTTTTTGAGC TGTGTAAGGA CCTTTCAATT CACTTTACAT TAGAATACGA
•	TATGTGATGG TAAGTATTAA CCCAGCTTCC TGAGTGATGG CGTGAGGGCG GAACCCAGGT
40	TCATGAAAAG TATTCCTATT ATTGAATTTT ACCAGTTATT TTCAGAAGTT TGATACCAT
	TGAGTTGATT TCATAACATG TCCTCTAACT GCACTTGATA GCAAATTATC TTGTTATCCT
	GAGTTGTAGC CAACAATGAC TTGGAGAATC TATGCAATAC TCAGTTTTAT TACTTCTAAG
	CTCATTTTGA AGATAATACT ACCCATGGCT GATTTGTTAC TATAAAATAG GTTTAGTATT
	TGCTGTCTGG AAACATTCTG ATTAGTGTCT CCTGGGAGGA TTATAAATTT ATAGTACCCA

	AAGANTAAAC NTGTTGTTTC CCTTTCCTAA	ACTTTTAGTG NATAATNCAG TCTCTGCCGT
	CTCTCATTTT CATCACTTGC CCTCCANAGC	TCCCATCTCA CTGAATTCTT GCAGTGTTCT
	CARTCATTGAG AGCCCCAANG TGGGTCTTAT	AACAGCCAGT CAGCAACATT TCTGTTTTTC
	ATCTGACACC AAGGGTCTCG TCTCTTTGCT	TTTCTACCAG TTATTCTGGG CTCTTCAGCT
5	CTARGARG TATAGGTCCT GARATCTTTC	CCTACCTTCT CAATTTCCTG GGGAGGGCTT
5	CTARACTURE THE TOTAL AND THE T	ATTTGAAGAT AATTGAGAAA TGAATGGAAA
	CTCANATTCA ACCOTTCATG TTAAGAGATT	GCAAGTTATG CTATCACCAA ATAGATTTTT
	MCMCCCMCAC ACCATTAATT CATACTCCAT	ATTATGTGTT GACTTTATCA TTGAGGTCCT
	COCACAMCAM ACCAMTGCCA TGATATAATT	TGAGCTACTG ATACTATAGT GTTGCTTCTG
1.0	CHCHTCHTAC CAGAACATCT AAAATATATT	AGGATTTTTT TAATGGCAGA GGAAATGAAG
10	ACMARATICA CATACTCCTT GTCCTAGTGA	A TTTACAGTTT AGCAAGACAT ACAAGCAAAA
	CAMURA A CURA A A GCATGATA A CTACTTTAA	A TAAAGCATTT TTTAAATTCA TTGGTAACAT
	ARCARCE AGARGETTT AGCAAACCCT	TCCCAAAAGA AATGATTGAC AAGTTATATG
	AGAMAAM TCACCCAAAC GAAATTCGAC	C TTTCTAAAGC CAAATTATTT GACATTGGTT
1 5		T CTCCCATGTT CTTTATTCTT TGACTGTTAT
15	ARCHARCATT TCTACAGAGA AATGGCAATT	T TCAAAACAGA GGGCTTTGAT GGATTAATTG
	CORRECT ATTC ATTCCTTCATC TACAGTATCT	T TGTCAGGTAC TTGAGAAAAT AATGTACTTA
	A A COURT COTO TOTAL COTTA COT	T TCTATACTGT AAGTTGGGGA AAAAAGTATT
	MUCHCATT COTATTGGG CTACTTGAA	A ATTCCCACCA ACTTTGCCAA TACCAGTGTT
20	CTCTATAACC CAGAATTCAG AATTAGCTG	C AATTAAGGGA ATTCACAGCT TTTCTGTAGT
20	CACACACAA ATTGAAATTA AAGAAAAAA	G AAATAGTGGG AGGACAAATG AGGTTTTACC
	TOTAL ACTUAL AND ACAGATT TAAGAACAA	G CCTTATACCT AGATTTATTA ATACTTTGAG
	CHATCACACC CAAGAGAAGC TTAGAAATA	C GGCAGAATGG GCTTTCTTTG TTCTTCTCCC
	ACCUARGO OF TOTAL TATAL TATAL	T TTTAAGAGAG AAGGGAAGTG TCTCTCCTGG
25	CMCACATTA TTAGGAAATA CAGAGTGTT	T TCATAATGCG TAAAGTCTAG TCCATTTAAG
25	TOTAL AND THE TATATTATT	TT GAGCAGGAAG GCAGAGACCT TAAACTGCTC
	CACCCAATTC ATTTACACA AAAGATTAA	AA AAGAAAAAAA CAGTGTCAAA AGGTCAAGTG
	CCCAGTGTTG CACAACTAAT GATAGGCAA	AA ACAAGAGAGG AGGGGAAAAA AAAGAATCCT
	CACTCCCGAC TCCTAGTTCA GTGTGTTCT	TT CACTGTTAGA AGGTGCTGCT GAACATAGTT
30	ATACCATATG AAGATCACAC TATCTATTT	TG AGATTGTAGA AATTTGATTA CTGCAGAGCT
30	CTGGCTGGGC TAGATTCACT TCTTATTCT	TT TATTGCATTG CAGTATTCTT AAGAGATAAA
	TEGETETTT AGAATCAGAT TEGECETTEG	GC TGTTGAAATG AGGCATTAAT TACCTTGGTA
	GCTGACACAT TCTTACAGGT CAGGGGCTT	TG ATGAAGTTTA TCTTCTTCCC TTGTTCCTGT
	GATTGCTCTG TAAATAGAAC ACATTCAGA	AG CCCTTGAATG CACTTGCAGC CTGTGCCTCC
35	CACAGTGATC GATGGTCAGA TAATGGGA	GT TTAATGACCA GTACTGAGAG AGATTATTTC
33	CATGGCTGCT ATGGGCCAAG AAAGCTGG	GT GGTCAGAAAG GGACCTTTTC CAGACTCTCC
ů.	TGGGTTGTGT TATTTCTTTG ACATTGGT	TT CCTTTCATGA GGCGCCAAGG TGTATTTGTA
	AGTTGTCCAG TGTTGCACAG CTAATGAG	GG GCAAAACAAG AAAGGAGGAA AAAAAAAGTA
	TCCTCACTCC TAGTTCAGTG TGTTCTTC	AC TGTTGGAAGG TACTGCTGAA CACAGTTATA
40	TGTTCAGTGT ACAATGTACA ATGTTCTT	TG ACATTCCGTC ATTTGAAGGA TGGGTCCTAT
- •	GGATAGTATC ATCTGCATGG TTTTGAAA	AC AAAAGATATA AGCTAATTTT GCCCTGTCTA
	GTGACTACGA GACAGGGAGA GAAAATCT	GA ATATTTGTTA AAGTAGACAC AGACCCATAA
	ATTGAAAAGG ACACTAATCC TGCCTTAG	GA GACAGTAAGG CACTTGTCCC TGCTATCTAT
	TAGCTGTGTG TCCTTGGACA GCTCATTG	GCT TCTTTCTGAG CCACCGTTAC ACGTTATCTA $3^{\prime\prime}$
		- 1

	TCAAGTGAGC AGGTTGTACA CTAGATGAAT TCACAGGTCC TTCTTCCAAG TGCTTTTCTA
	ATCTTCATAA TTTAGATAAT CTCTCAGTAG CAAAACAGTG TACAATATGG TCAATCTGAG
	ATCTTCATAA TITAGATAAT CICICAGTAG CHULLICAGTAG CHULLICAGTA
	ATTTTTAGGG GGAGAATTTT AGGGAACATC ACIDITIOGEN CONTROL CONTROL CATGGTAGAA GGGATGAGAG TTCTCTCTGG
_	AGCCTTTTGT ATTAGGTCAC TAATCCCAAA GGCCCCACCT CCTAAGACCA TCACTTTTGG
5	AGCCTTTTGT ATTAGGTCAC TAATCCCAAA GGCCCCACCT COTTAGATCATTG CATTAGAGAT GATTGGGGTT TTAACATGAT TTTAGAGGGA CATAAACCTT CATCTCATTG CATTAGAGAT
	ATTTGAAAGC TCAGCTCACG TGTATTCCTC CACAGCTCAC ATGTATTCCT CCATCACCCA
	ACCTGATGGC TTTGAACGTT GTAGACATAA ATCCTTTCAT CATTATCAAG AATATTGCCA
	AAAGCTTCTC AGAAATTATG AGGGGTTTTT TTAGTTTCTA AAATATTCTC AAAGAAAGTC
10	CCATGTACTA ATGTTTGCCT TTTGATGAAA AAGGATGAAA TCTTAATGAT TGCCTTAATA
	AGCTCAACAA TGCTTGTTAG TTGAGTCTTC TTATTGTGCT GATTCTTATA AACAACAACA
	TTCAGTATAA ACATTAATGT ATGTGATTCA CTAAGGTTTT TGCATGATTC TCTGTGAGGC
	TTCAGATGTC TCTTGGATTA TGTGTCTTTT TTTCATCGCC AGCATCCTGT GGCAGCTGCT
	ACTCATTTGC TTCTATGGGT ATGCTAGAAG CGAGAATCCG TATACTAACC AACAATTCTC
15	AGACCCCAAT CCTAAGCCCT CAGGAGGTTG TGTCTTGTAG CCAGTATGCT CAAGGTAAGT
	GTTGCATTTC AGACACCATT TATGAGCTAT TTACCTGTGT GCAGCTGGCT GTTGTTGGCA
	AAGGCAAAAG GATGATGCAG TAGAGAGAGC GCAGTGTCTA TAGTCAGAAA ATCTGAGTGC
	AAGTCTGGCC CTATCACTTA TTAATGGATG ATTGCTCATG GAATTTACTG TACCATCCAG
	CAAAATGTCA ATAGTTACTA TATATTGAGT GAGCTCTGCT TGTTACATAT ATTGCCTAAC
20	AATGCTCAAA ACTCTGAGAA GTAGTAAGTA TAATCCCTAT TTATGGGCGG GGAACAGGAA
	CTAAGAAATT TTTCTAATAA TTTGAAGGTC TCACAGCTTT TAGCATTGGA GTTTCACTTC
	TAATCATCGT CTCCAAAACC CAACTTTTAT TAAAACTATA CTAACACTGG TTTCTCTCTG
	GGAGAATTTT AAAATTCTGT ACTTAGGGCT GGGCACAGTG GCTTATGCCT ATAACCCTAT
	CACTTTGGGA GGCTGAGATG GGTGAATCTC TTGAGTCCTA GAGTTTGAGA CCAGCCTGGG
25	CAACACGGCG AAACCCCTTC TCTATTAAAA ATACAAAAAA TTAGCTGGGC GTGGTGGTGT
	GTGCTTGTAG TCCCAGCTAT TCAGGAGGCG GAGGTGTAAG AATCACCTGA GCCCAGGAGG
	TCAAGGCTGC AGTGAGCCGA TATCATGCCA CTGCACTCCA GCCTGGGCAA ACGGAGTGAG
	GCCCTGTTAT GAAAAAAAA AAATCTGTAC TTAGGCTTTC AGATCAGGCT GTATGTGATG
	TATGTCGAAA ACACAGCTAT AATTGATTGA GGGAGAAACG TTACCATTTT AAAGTTTATG
30	CTTTCAAGCC CAGATTTGGC CACTAGGAAT TTCCCAGCTC ACTAGTGAAA CTGCTGATGA
	GTGATTATTT GCCAGTGAGC CTTTCATTCT TTCTAAAATA TGTACTACTA GTTGTGACTT
	GTAGGCTATA GGGGCTATAA TATATCAAGA CAATCTTTAT CCTCATGAAG CTTACAGTTA
	AGTAAGAGAT AGAGATTAAA TAATTATAAC AACAGAGTGA AGAACAGTGA AGAAAAAGTA
	CAGAGTTATA ATATATAA TAGGGCCAGG ACTGCATGAG GAAGGTAGGA AAGACATTTC
35	GGCAAGAGGT TGTCAGGGAA AAGACTTGCT TGAGAAAGAG CCAAGTTGTG GGGTCTGGCT
33	GCTTAGCAAT GACCATAATA CCTAACTTTT GCTATTTTTA CATGAAGTAA CTAATTTAAC
-	CCTATGAGGA AAGTACTACT ACCATCTAGA TTTTACAGGT AAGTAAGCAG AGATACAGAG
	AAGTTAAACT CTTCACACGG CTTTGGCTTT AAACCTATAT AGGCTTCAGA GCCTCCCCAC
	TTAACCACTT TGCCATAGCT ACATCCATAT TAGGTGCTAA GTAGATATCT GTTAAGTAGA
40	AGGAGGATGA AAGGATAGTT AGCTAGTTGG AAAATGGATG GATGAATGAA GTGATGCTTA
<b>-</b> ± ∪	AGCTAAGAAC AACTTTCAGG GGTAACATGC AAAGAATAAT GGAGCAAAGA AGAAAAAATA
	GAAAATGGGA TAATCCTTTT TCTACTAAGG GGTAACCATG TGTGTTATTC ATCTTCAGGC
	TGTGAAGGCG GCTTCCCATA CCTTATTGCA GGAAAGTACG CCCAAGATTT TGGGCTGGTG
	GAAGAAGCTT GCTTCCCCTA CACAGGCACT GATTCTCCAT GCAAAATGAA GGAAGACTGC
	3.5

10031569.012502

PCT/US00/20400 WO 01/07663

TTTCGTTATT	ACTCCTCTGA	GTACCACTAT	GTAGGAGGTT	TCTATGGAGG	CTGCAATGAA
				CAGTTGCTTT	
				CTGGTCTAAG	
				GCTATGGCAC	
				CCGGCTGGGG	
				AGAGCATAGC	
				CATAATGATC	
				GCAGCAATCT	
				ACTGCCCTTA	
				AATTAAGTGG	
	AGCTATGAAG			TTTGTAATTC	
TATTTTTAA	AATCAATGTG	AAAACATAGA	CTTATTTTA	AATTGTACCA	ATCACAAGAA
AATAATGGCA	ATAATTATCA	AAACTTTTAA	AATAGATGCT	CATATTTTTA	AAATAAAGTT
TTAAAA					

15

5

10

The corresponding cDNA sequence for CTSC is provided below (SEQ ID NO: 2):

	_		SEG ID NO				
						tgggccctc c	
20	61	gccgccctcc	tgctgcttct	ctccggcgac	ggcgccgtgc	gctgcgacac	acctgccaac
	121	tgcacctatc	ttgacctgct	gggcacctgg	gtcttccagg	tgggctccag	cggttcccag
	181	cgcgatgtca	actgctcggt	tatgggacca	caagaaaaaa	aagtagtggt	gtaccttcag
	241	aagctggata	cagcatatga	tgaccttggc	aattctggcc	atttcaccat	catttacaac
	301	caaggctttg	agattgtgtt	gaatgactac	aagtggtttg	ccttttttaa	gtataaagaa
25	361	gagggcagca	aggtgaccac	ttactgcaac	gagacaatga	ctgggtgggt	gcatgatgtg
	421	ttgggccgga	actgggcttg	tttcaccgga	aagaaggtgg	gaactgcctc	tgagaatgtg
	481	tatgtcaaca	cagcacacct	taagaattct	caggaaaagt	attctaatag	gctctacaag
						agtcttggac	
						ggagaagtgg	
30						aaatacagca	
	721	catttgccaa	catcttggga	ctggagaaat	gttcatggta	tcaattttgt	cagtcctgtt
	781	cgaaaccaag	catcctgtgg	cagctgctac	tcatttgctt	ctatgggtat	gctagaagcg
						taagccctca	
						cataccttat	
35	961	tacgcccaag	attttgggct	ggtggaagaa	gcttgcttcc	cctacacagg	cactgattct
						ctgagtacca	
	1081	ggtttctatg	gaggctgcaa	tgaagccctg	atgaagcttg	agttggtcca	tcatgggccc
	1141	atggcagttg	cttttgaagt	atatgatgac	ttcctccact	acaaaaaggg	gatctaccac
	1201	. cacactggtc	taagagaccc	tttcaacccc	tttgagctga	ctaatcatgc	tgttctgctt
40	1261	gtgggctatg	gcactgacto	agcctctggg	atggattact	. ggattgttaa	aaacagctgg
	1321	ggcaccggct	ggggtgagaa	tggctacttc	cggatccgca	gaggaactga	tgagtgtgca
	1381	l attgagagca	tagcagtggc	agccacacca	attcctaaat	: tgtagggtat	gccttccagt
	1443	l atttcataat	gatctgcato	agttgtaaag 36	gggaattggt	atattcacag	actgtagact

1501 ttcagcagca atctcagaag cttacaaata gatttccatg aagatatttg tcttcagaat
1561 taaaactgcc cttaatttta atataccttt caatcggcca ctggccattt ttttctaagt
1621 attcaattaa gtgggaattt tctggaagat ggtcagctat gaagtaatag agtttgctta
1681 atcatttgta attcaaacat gctatatttt ttaaaatcaa tgtgaaaaca tagacttatt
1741 tttaaattgt accaatcaca agaaaataat ggcaataatt atcaaaactt ttaaaataga
1801 tgctcatatt tttaaaataa agttttaaaa ataactgc

The wild type CTSC protein sequence is set forth below as SEQ ID NO: 3:

10 MGAGPSLLLAALLLLLSGDGAVRCDTPANCTYLDLLGTWVFQVG
SSGSQRDVNCSVMGPQEKKVVVYLQKLDTAYDDLGNSGHFTIIYNQGFEIVLNDYKWF
AFFKYKEEGSKVTTYCNETMTGWVHDVLGRNWACFTGKKVGTASENVYVNTAHLKNSQ
EKYSNRLYKYDHNFVKAINAIQKSWTATTYMEYETLTLGDMIRRSGGHSRKIPRPKPA
PLTAEIQQKILHLPTSWDWRNVHGINFVSPVRNQASCGSCYSFASMGMLEARIRILTN
NSQTPILSPQEVVSCSQYAQGCEGGFPYLIAGKYAQDFGLVEEACFPYTGTDSPCKMK
EDCFRYYSSEYHYVGGFYGGCNEALMKLELVHHGPMAVAFEVYDDFLHYKKGIYHHTG
LRDPFNPFELTNHAVLLVGYGTDSASGMDYWIVKNSWGTGWGENGYFRIRRGTDECAI
ESIAVAATPIPKL.

Papillon Lefevre syndrome (PLS) is an autosomal recessive disorder characterized by palmoplantar hyperkeratosis and severe early onset periodontitis that results in the premature loss of the primary and secondary dentitions. The 46 kb CTSC gene consists of 7 exons and is mutated in PLS patients. Sequence analysis of CTSC from PLS affected individuals from thirty-two Turkish families identified four different mutations. An exon 6 nonsense mutation (856C->T) introduces a premature stop codon at amino acid 286. Three exon 2 mutations were identified including a single nucleotide deletion (1047delA) of codon 349 introducing a frameshift and premature termination codon, a two base pair deletion (1028-1029delCT) that results in introduction of a stop codon at amino acid 343, and a G->A substitution in codon 429 (1286G->A) introducing a premature termination codon. All PLS affected individuals examined were homozygous for cathepsin C mutations inherited from a common ancestor. Parents and siblings heterozygous for cathepsin C mutations do not show either the palmoplantar hyperkeratosis or severe early onset periodontitis characteristic of PLS.

40

5

20

25

30

addition to the 5 families described above, Table I summarizes CTSC mutations identified in 27 other families presenting with symptoms of PLS.

Haim-Munk syndrome is a rare condition associated with congenital palmoplantar keratosis, pes planus, onychogyrphosis, periodontosis, arachnodactyly and acroosteolysis. In an additional embodiment of the invention, a mutation in cathepsin C causing Haim-Munk Syndrome has been identified. It appears that substitution of an A for a G at CTSC nucleotide position 857 in Exon 6 is responsible for this syndrome in patients.

Based on the data presented herein, it appears that additional mutations or functional polymorphisms are associated with other pathological conditions, including, but not limited to prepubertal periodontitis (PPP), early onset periodontal disease or other forms of gum disease. For example, as shown herein, PPP is caused a substitution of a G for an A at position 1040 in the CTSC coding sequence. Thus, the invention also provides methods for screening the CTSC gene for alterations associated with these disease states.

# I. <u>Preparation of Altered Human CTSC-Encoding Nucleic</u> <u>Acid Molecules, CTSC Proteins, and Antibodies</u> <u>Thereto</u>

#### A. Nucleic Acid Molecules

Nucleic acid molecules encoding the human CTSC proteins of the invention may be prepared by two general methods: (1) synthesis from appropriate nucleotide triphosphates, or (2) isolation from biological sources. Both methods utilize protocols well known in the art. The availability of nucleotide sequence information, such as a DNA having the sequence of SEQ ID NOS:1-2 enables preparation of an isolated nucleic acid molecule of the invention by oligonucleotide synthesis. Synthetic oligonucleotides may be prepared by the

35

30

5

10

15

20

PCT/US00/20400

WO 01/07663

5

10

15

20

25

30

phosphoramidite method employed in the Applied Biosystems 38A DNA Synthesizer or similar devices. resultant construct may be purified according to methods known in the art, such as high performance liquid chromatography (HPLC). Long, double-stranded polynucleotides, such as a DNA molecule of the present invention, must be synthesized in stages, due to the size limitations inherent in current oligonucleotide Thus, for example, a 4.7 kb doublesynthetic methods. stranded molecule may be synthesized as several smaller segments of appropriate complementarity. Complementary segments thus produced may be annealed such that each segment possesses appropriate cohesive termini for attachment of an adjacent segment. Adjacent segments may be ligated by annealing cohesive termini in the presence of DNA ligase to construct an entire 4.7 kb double-stranded molecule. A synthetic DNA molecule so constructed may then be cloned and amplified in an appropriate vector.

Nucleic acid sequences encoding the altered human CTSC proteins of the invention may be isolated from appropriate biological sources using methods known in In a preferred embodiment, a cDNA clone is isolated from a cDNA expression library of human origin. In an alternative embodiment, utilizing the sequence information provided by the cDNA sequence, human genomic clones encoding altered CTSC proteins may be isolated.

Table 1 sets forth several different mutations associated with particular PPKs and PPP. Altered CTSCspecific probes for identifying such sequences may be between 15 and 40 nucleotides in length. For probes longer than those shown above, the additional contiguous nucleotides are provided within SEQ ID NOS:1 and 2.

Additionally, cDNA or genomic clones having homology with human CTSC may be isolated from other species using oligonucleotide probes corresponding to predetermined sequences within the human CTSC encoding

nucleic acids.

5

10

15

20

25

In accordance with the present invention, nucleic acids having the appropriate level of sequence homology with the protein coding region of SEQ ID NO:1 may be identified by using hybridization and washing conditions of appropriate stringency. For example, hybridizations may be performed, according to the method of Sambrook et al., Molecular Cloning, Cold Spring Harbor Laboratory (1989), using a hybridization solution comprising: 5X SSC, 5X Denhardt's reagent, 1.0% SDS, 100 µg/ml denatured, fragmented salmon sperm DNA, 0.05% sodium pyrophosphate and up to 50% formamide. Hybridization is carried out at 37-42°C for at least six hours.

Following hybridization, filters are washed as follows: (1) 5 minutes at room temperature in 2X SSC and 1% SDS; (2) 15 minutes at room temperature in 2X SSC and 0.1%

(2) 15 minutes at room temperature in 2X SSC and 0.1% SDS; (3) 30 minutes-1 hour at 37°C in 1X SSC and 1% SDS;

(4) 2 hours at 42-65 oC in 1X SSC and 1% SDS, changing the solution every 30 minutes.

Nucleic acids of the present invention may be maintained as DNA in any convenient cloning vector. In a preferred embodiment, clones are maintained in a plasmid cloning/expression vector, such as pBluescript (Stratagene, La Jolla, CA), which is propagated in a suitable *E. coli* host cell.

Altered CTSC-encoding nucleic acid molecules of the invention include cDNA, genomic DNA, RNA, and fragments thereof which may be single- or double-stranded. Thus, this invention provides oligonucleotides having sequences capable of hybridizing with at least one sequence of a nucleic acid molecule of the present invention, such as selected segments of the DNA having SEQ ID NO:1. Also contemplated in the scope of the present invention are oligonucleotide probes which specifically hybridize with the mutated CTSC genes of the invention while not hybridizing with the wild type sequence under high stringency conditions. Primers

35

capable of specifically amplifying the altered CTSC encoding nucleic acids described herein are also contemplated herein. As mentioned previously, such oligonucleotides are useful as probes and primers for detecting, isolating or amplifying altered CTSC genes.

Antisense nucleic acid molecules may be targeted to translation initiation sites and/or splice sites to inhibit the expression of the CTSC gene or production of the CTSC protein of the invention. Such antisense molecules are typically between 15 and 30 nucleotides in length and often span the translational start site of CTSC encoding mRNA molecules.

Alternatively, antisense constructs may be generated which contain the entire CTSC cDNA in reverse orientation. Such antisense constructs are easily prepared by one of ordinary skill in the art.

It will be appreciated by persons skilled in the art that variants (e.g., allelic variants) of CTSC sequences exist in the human population, and must be taken into account when designing and/or utilizing oligonucleotides of the invention. Accordingly, it is within the scope of the present invention to encompass such variants, with respect to the CTSC sequences disclosed herein or the oligonucleotides targeted to specific locations on the respective genes or RNA transcripts. Accordingly, the term "natural allelic variants" is used herein to refer to various specific nucleotide sequences of the invention and variants thereof that would occur in a human population. usage of different wobble codons and genetic polymorphisms which give rise to conservative or neutral amino acid substitutions in the encoded protein are Such variants would not examples of such variants. demonstrate altered CTSC activity. Additionally, the term "substantially complementary" refers to oligonucleotide sequences that may not be perfectly matched to a target sequence, but such mismatches do not

35

30

5

10

15

20

PCT/US00/20400

WO 01/07663

materially affect the ability of the oligonucleotide to hybridize with its target sequence under the conditions described.

5

10

15

20

25

#### B. Proteins

Full-length, altered, human CTSC proteins of the present invention may be prepared in a variety of ways, according to known methods. The proteins may be purified from appropriate sources, e.g., transformed bacterial or animal cultured cells or tissues, by immunoaffinity purification. However, this is not a preferred method due to the low amount of protein likely to be present in a given cell type at any time. availability of nucleic acid molecules encoding CTSC protein enables production of the protein using in vitro expression methods known in the art. For example, a cDNA or gene may be cloned into an appropriate in vitro transcription vector, such as pSP64 or pSP65 for in vitro transcription, followed by cell-free translation in a suitable cell-free translation system, such as wheat germ or rabbit reticulocyte lysates. transcription and translation systems are commercially available, e.g., from Promega Biotech, Madison, Wisconsin or Gibco-BRL, Gaithersburg, Maryland.

Alternatively, according to a preferred embodiment, larger quantities of CTSC protein may be produced by expression in a suitable prokaryotic or eukaryotic system. For example, part or all of a DNA molecule, such as a DNA having SEQ ID NOS:1 or 2 containing an alteration set forth in Table 1 may be inserted into a plasmid vector adapted for expression in a bacterial cell, such as E. coli. Such vectors comprise the regulatory elements necessary for expression of the DNA in the host cell positioned in such a manner as to permit expression of the DNA in the host cell. Such regulatory elements required for expression include

35

promoter sequences, transcription initiation sequences and, optionally, enhancer sequences.

The human CTSC protein produced by gene expression in a recombinant procaryotic or eukaryotic system may be purified according to methods known in the art. preferred embodiment, a commercially available expression/secretion system can be used, whereby the recombinant protein is expressed and thereafter secreted from the host cell, and readily purified from the surrounding medium. If expression/secretion vectors are not used, an alternative approach involves purifying the recombinant protein by affinity separation, such as by immunological interaction with antibodies that bind specifically to the recombinant protein or nickel columns for isolation of recombinant proteins tagged with 6-8 histidine residues at their N-terminus or Cterminus. Alternative tags may comprise the FLAG epitope or the hemagglutinin epitope. Such methods are commonly used by skilled practitioners.

The human CTSC protein of the invention, prepared by the aforementioned methods, may be analyzed according to standard procedures. For example, such protein may be subjected to amino acid sequence analysis, according to known methods.

The present invention also provides antibodies capable of immunospecifically binding to proteins of the invention. Polyclonal antibodies directed toward altered human CTSC proteins may be prepared according to standard methods. In a preferred embodiment, monoclonal antibodies are prepared, which react immunospecifically with the various epitopes of the CTSC protein described herein. Monoclonal antibodies may be prepared according to general methods of Köhler and Milstein, following standard protocols. Polyclonal or monoclonal antibodies that immunospecifically interact with altered CTSC proteins can be utilized for identifying and purifying such proteins. For example, antibodies may be utilized

20

5

10

15

25

35

for affinity separation of proteins with which they immunospecifically interact. Antibodies may also be used to immunoprecipitate proteins from a sample containing a mixture of proteins and other biological molecules. Other uses of anti-CTSC antibodies are described below.

# II. DETECTION OF KERATODERMAL DISORDERS/DYSPLASIAS and PERIODONTAL DISEASE-ASSOCIATED MUTATIONS AND DIAGNOSTIC SCREENING ASSAYS THEREFORE

Currently, the most direct method for mutational analysis is DNA sequencing, however it is also the most labor intensive and expensive. It is usually not practical to sequence all potentially relevant regions of every experimental sample. Instead some type of preliminary screening method is commonly used to identify and target for sequencing only those samples that contain mutations. Single stranded conformational polymorphism (SSCP) is a widely used screening method based on mobility differences between single-stranded wild type and mutant sequences on native polyacrylamide gels. Other methods are based on mobility differences in wild type/mutant heteroduplexes (compared to control homoduplexes) on native gels (heteroduplex analysis) or denaturing gels (denaturing gradient gel electrophoresis). Sample preparation is relatively easy in these assays, and conditions for electrophoresis required to generate the often subtle mobility differences that form the basis for identifying the targets that contain mutations are known to those of skill in the art. Another parameter to be considered is the size of the target region being screened. general, SSCP is used to screen target regions no longer than about 200-300 bases.

Another type of screening technique currently in use is based on cleavage of unpaired bases in

44

5

10

20

15

25

30

10031889 O1E50E

WO 01/07663 PCT/US00/20400

heteroduplexes formed between wild type probes hybridized to experimental targets containing point The cleavage products are also analyzed by mutations. gel electrophoresis, as subfragments generated by cleavage of the probe at a mismatch generally differ significantly in size from full length, uncleaved probe and are easily detected with a standard gel system. Mismatch cleavage has been effected either chemically (osmium tetroxide, hydroxylamine) or with a less toxic, enzymatic alternative, using RNase A. The RNase A cleavage assay has also been used, although much less frequently, to screen for mutations in endogenous mRNA targets for detecting mutations in DNA targets amplified by PCR. A mutation detection rate of over 50% was reported for the original RNase screening method.

10

15

20

25

30

35

A newer method to detect mutations in DNA relies on DNA ligase which covalently joins two adjacent oligonucleotides which are hybridized on a complementary target nucleic acid. The mismatch must occur at the site of ligation. As with other methods that rely on oligonucleotides, salt concentration and temperature at hybridization are crucial. Another consideration is the amount of enzyme added relative to the DNA concentration. In summary, exemplary approaches for detecting alterations in CTSC encoding nucleic acids or polypeptides/proteins include:

- a) comparing the sequence of nucleic acid in the sample with the wild-type CTSC nucleic acid sequence to determine whether the sample from the patient contains mutations; or
- b) determining the presence, in a sample from a patient, of the polypeptide encoded by the CTSC gene and, if present, determining whether the polypeptide is full length, and/or is mutated, and/or is expressed at the normal level; or
- c) using DNA restriction mapping to compare the restriction pattern produced when a restriction enzyme

cuts a sample of nucleic acid from the patient with the restriction pattern obtained from normal CTSC gene or from known mutations thereof; or,

- d) using a specific binding member capable of binding to a CTSC nucleic acid sequence (either normal sequence or known mutated sequence), the specific binding member comprising nucleic acid hybridizable with the CTSC sequence, or substances comprising an antibody domain with specificity for a native or mutated CTSC nucleic acid sequence or the polypeptide encoded by it, the specific binding member being labeled so that binding of the specific binding member to its binding partner is detectable; or,
- e) using PCR involving one or more primers based on normal or mutated CTSC gene sequence to screen for normal or mutant CTSC gene in a sample from a patient.

A "specific binding pair" comprises a specific binding member (sbm) and a binding partner (bp) which have a particular specificity for each other and which in normal conditions bind to each other in preference to other molecules. Examples of specific binding pairs are antigens and antibodies, ligands and receptors and complementary nucleotide sequences. The skilled person is aware of many other examples and they do not need to be listed here. Further, the term "specific binding pair" is also applicable where either or both of the specific binding member and the binding partner comprise a part of a large molecule. In embodiments in which the specific binding pair are nucleic acid sequences, they will be of a length to hybridize to each other under conditions of the assay, preferably greater than 10 nucleotides long, more preferably greater than 15 or 20 nucleotides long.

In most embodiments for screening for susceptibility alleles, the CTSC nucleic acid in the sample will initially be amplified, e.g. using PCR, to increase the amount of the analyte as compared to other  $4\ell$ 

35

30

10

15

20

10031<u>9</u>69.01E50E

WO 01/07663 PCT/US00/20400

sequences present in the sample. This allows the target CTSC sequences to be detected with a high degree of sensitivity if they are present in the sample. This initial step may be avoided by using highly sensitive array techniques that are becoming increasingly important in the art.

The identification of the CTSC gene and its association with keratodermal disorders/dysplasias and peridontal diseases paves the way for aspects of the present invention to provide the use of materials and methods, such as are disclosed and discussed above, for establishing the presence or absence in a test sample of a variant form of the gene, in particular an allele or variant specifically associated with PLS, HMS or periodontal diseases. This may be for diagnosing a predisposition of an individual to PLS, HMS or periodontal disease. It may be for diagnosing PLS, HMS or periodontal disease in a patient with the disease as being associated with the altered CTSC gene.

This allows for planning of appropriate therapeutic and/or prophylactic measures, permitting stream-lining of diagnosis, treatment and outcome assessments. The approach further stream-lines treatment by targeting those patients most likely to benefit.

According to another aspect of the invention, methods of screening drugs for therapy to identify suitable drugs for restoring CTSC product functions are provided.

The CTSC polypeptide or fragment employed in drug screening assays may either be free in solution, such as gingival crevicular fluid, affixed to a solid support or within a cell. One method of drug screening utilizes eukaryotic or prokaryotic host cells which are stably transformed with recombinant polynucleotides expressing the polypeptide or fragment, preferably in competitive binding assays. Such cells, either in viable or fixed form, can be used for standard binding assays. One may

30

10

15

20

25

WO 01/07663

5

10

15

20

25

determine, for example, formation of complexes between a CTSC polypeptide or fragment and the agent being tested, or examine the degree to which the formation of a complex between a CTSC polypeptide or fragment and a known ligand is interfered with by the agent being tested.

Another technique for drug screening provides high throughput screening for compounds having suitable binding affinity to CTSC polypeptides and is described in detail in Geysen, PCT published application WO 84/03564, published on Sep. 13, 1984. Briefly stated, large numbers of different, small peptide test compounds are synthesized on a solid substrate, such as plastic pins or some other surface. The peptide test compounds are reacted with CTSC polypeptide and washed. CTSC polypeptide is then detected by methods well known in the art.

A further technique for drug screening involves the use of host eukaryotic cell lines or cells (such as described above) which have a nonfunctional CTSC gene. These host cell lines or cells are defective at the CTSC polypeptide level. The host cell lines or cells are grown in the presence of drug compound. The rate of growth of the host cells is measured to determine if the compound is capable of regulating the growth of CTSC defective cells.

The goal of rational drug design is to produce structural analogs of biologically active polypeptides of interest or of small molecules with which they interact (e.g., agonists, antagonists, inhibitors) in order to fashion drugs which are, for example, more active or stable forms of the polypeptide, or which, e.g., enhance or interfere with the function of a polypeptide in vivo. See, e.g., Hodgson, (1991) In one approach, one first Bio/Technology 9:19-21. determines the three-dimensional structure of a protein polypeptide) or, for example, of interest (e.g., CTSC

35

of the CTSC-substrate complex, by x-ray crystallography, by nuclear magnetic resonance, by computer modeling or most typically, by a combination of approaches. Less often, useful information regarding the structure of a polypeptide may be gained by modeling based on the structure of homologous proteins. An example of rational drug design is the development of HIV protease inhibitors (Erickson et al., (1990) Science 249:527-533). In addition, peptides (e.g., CTSC polypeptide) may be analyzed by an alanine scan (Wells, 1991) Meth. Enzym. 202:390-411. In this technique, an amino acid residue is replaced by Ala, and its effect on the peptide's activity is determined. Each of the amino acid residues of the peptide is analyzed in this manner to determine the important regions of the peptide.

It is also possible to isolate a target-specific antibody, selected by a functional assay, and then to solve its crystal structure. In principle, this approach yields a pharmacore upon which subsequent drug design can be based. It is possible to bypass protein crystallography altogether by generating anti-idiotypic antibodies (anti-ids) to a functional, pharmacologically active antibody. As a mirror image of a mirror image, the binding site of the anti-ids would be expected to be an analog of the original molecule. The anti-id could then be used to identify and isolate peptides from banks of chemically or biologically produced banks of peptides. Selected peptides would then act as the pharmacore.

Thus, one may design drugs which have, e.g., improved CTSC polypeptide activity or stability or which act as inhibitors, agonists, antagonists, etc. of CTSC polypeptide activity. By virtue of the availability of cloned CTSC sequences, sufficient amounts of the CTSC polypeptide may be made available to perform such analytical studies as x-ray crystallography. In addition, the knowledge of the CTSC protein sequence

provided herein will guide those employing computer modeling techniques in place of, or in addition to x-ray crystallography.

III Therapeutics

# A. Pharmaceuticals and Peptide Therapies

The discovery that mutations in the CTSC gene give rise to PLS, HMS, and periodontal disease facilitates the development of pharmaceutical compositions useful for treatment and diagnosis of these syndromes and conditions. These compositions may comprise, in addition to one of the above substances, a pharmaceutcally acceptable excipient, carrier, buffer, stabilizer or other materials well known to those skilled in the art. Such materials should be non-toxic and should not interfere with the efficacy of the active ingredient. The precise nature of the carrier or other material may depend on the route of administration, e.g. oral, intravenous, cutaneous or subcutaneous, nasal, intramuscular, intraperitoneal routes.

Whether it is a polypeptide, antibody, peptide, nucleic acid molecule, small molecule or other pharmaceutically useful compound according to the present invention that is to be given to an individual, administration is preferably in a "prophylactically effective amount" or a "therapeutically effective amount" (as the case may be, although prophylaxis may be considered therapy), this being sufficient to show benefit to the individual.

#### B. Methods of Gene Therapy

As a further alternative, the nucleic acid encoding the authentic biologically active CTSC polypeptide could be used in a method of gene therapy, to treat a patient who is unable to synthesize the active "normal" polypeptide or unable to synthesize it at the normal

35

30

5

10

15

20

level, thereby providing the effect elicited by wildtype CTSC and suppressing the occurrence of "abnormal" CTSC associated with keratodermal disorders and dysplasias.

5

10

15

20

25

30

Vectors, such as viral vectors have been used in the prior art to introduce genes into a wide variety of different target cells. Typically the vectors are exposed to the target cells so that transformation can take place in a sufficient proportion of the cells to provide a useful therapeutic or prophylactic effect from the expression of the desired polypeptide. The transfected nucleic acid may be permanently incorporated into the genome of each of the targeted cells, providing long lasting effect, or alternatively the treatment may have to be repeated periodically.

A variety of vectors, both viral vectors and plasmid vectors are known in the art, see US Patent No. 5,252,479 and WO 93/07282. In particular, a number of viruses have been used as gene transfer vectors, including papovaviruses, such as SV40, vaccinia virus, herpes viruses including HSV and EBV, and retroviruses. Many gene therapy protocols in the prior art have employed disabled murine retroviruses.

Gene transfer techniques which selectively target the CTSC nucleic acid to oral tissues are preferred. Examples of this include receptor-mediated gene transfer, in which the nucleic acid is linked to a protein ligand via polylysine, with the ligand being specific for a receptor present on the surface of the target cells.

The following methods are provided to facilitate the practice of the present invention.

# 35 Family material and clinical diagnosis.

Five Turkish families were described previously [8]. All available family members provided consent for

PCT/US00/20400

WO 01/07663

5

10

15

20

25

30

35

the study and were clinically examined. A diagnosis of PLS was made in individuals with severe early onset periodontitis and the clinical appearance of hyperkeratosis on the palmar and plantar surfaces. All affected individuals also had hyperkeratosis on the knees. DNA was isolated from peripheral blood samples from all available members from these nuclear families using standard techniques (Qiamp Blood Kit, Qiagen).

# RNA Isolation, Amplification, and Tissue Expression Analysis.

Total RNA was generated from fresh tissue samples (gingiva, palm, sole, knee) using TRIZOL reagent (Molecular Research Center, Inc.; Cincinnati, OH) according to the manufacturer's protocol. To determine if cathepsin C was expressed in a given tissue, singletube RT-PCR was carried out using the Access RT-PCR System (Promega; Madison, WI), following the manufacturer's protocol. A portion of each reaction was visualized following agarose gel electrophoresis in the presence of ethidium bromide. Amplification primers located within exon 6  $\mathbf{F}$  5'-AGGAGGTTGTGTCTTGTAGCC-3'(nt. 857-877; SEQ ID NO: 4) and exon 7  $\mathbf{R}$  5'-AGTGCCTGTGTAGGGGAAGC-3' (NT 981-962; SEQ ID NO: 5) produce an amplicon of 123 base pairs from cDNA. A standard PCR protocol was followed with an annealing temperature of 65°C.

### GenBank accession numbers.

Full-length cDNA of CTSC (NM-001814) and full-length genomic DNA of CTSC contained within a BAC vector, Genbank accession number (AC011088). See SEQ ID NO: 1.

## Cathepsin C Activity Assay

In unafffected non-carriers, cathepsin C activity ranges from  $600-1200~\mu\text{mol/min/mg}$ . As carriers of a

PCT/US00/20400 WO 01/07663

cathepsin C mutation do not have clinical manifestations, measurement of cathepsin C enzymatic activity can be used to determine whether at-risk family members are carriers. Enzymatic activity can also be used to determine if individuals marrying into a family Carriers typically have approximately 50% are carriers. of normal enzyme activity. Determination of enzymatic activity can also be used to establish a diagnosis of PLS when mutational studies of cathepsin C have been negative. This is important in assuring that a diagnosis of PLS has been properly given to an individual with clinical symptoms suggestive of PLS.

Viable leukocyte pellets are obtained from lithium heparinized whole blood by mixing blood with 3 volumes of 3% dextran in normal saline, and allowing the red cells to settle for 45 min at room temperature. Cells are pelleted by centrifugation at 1500 rpm for 5 min at 4\*C. After washing and removal of contaminating red cells, leukocyte pellets are resuspended in dH20 and sonicated on ice for 5 sec each for total of 6 blasts using a Sonic 300 Dismembrator. Protein concentration is determined by the Lowry method.

Enzymatic activity is determined by measuring hydrolysis of the synthetic substrate glycyl-L-arginine-7-amido-4-methylcoumarin at a final concentration of 5 mM using a modified method. reactions are performed in duplicate. Twenty µl of leukocyte lysate are added to 200  $\mu l$  of  $Na_3PO_4$ buffer (0.1M, pH 6.5) in a 96 well plate and then substrate added. Reactions are allowed to proceed for 1 hr at room temperature at which time 10  $\mu l$  of glycine-NaOH buffer (0.5M, pH 9.8) is added to stop the reaction. Fluorescence is determined using a Perkin-Elmer LS50B luminescence spectrometer at 370-nm The amount of NHMec excitation and 460-nm emission. released is determined by generating a standard curve using NHMec. Cathepsin C activity is reported as µmol

35

30

5

10

15

20

5

10

15

20

25

NHMec released per min per mg protein.

### Sequencing and mutation analysis.

PCR primers were designed to cover the entire cathepsin C gene in overlapping fragments, from 955 nucleotides 5' to the start codon to 240 nucleotides 3' to the termination codon using cathepsin C (DPP-I) sequence data (Accession # U79415; SEQ ID NO: 1). The PCR products were prepared for sequencing by excising the bands from the agarose gel and extracting the fragments using a Qiagen Gel Clean-up Kit. The sense and antisense strand of each PCR product were directly sequenced on an ABI Prism 310 Genetic Analyzer (Perkin-Elmer) using four dye terminator chemistry. Approximately 1-3 ng of purified product and 3.2 pmol primer were added to premixed reagents from the ABI Prism Big Dye Terminator Cycle Sequencing Ready Reaction Kit, FS (Perkin-Elmer) and underwent a cycle sequencing reaction in a GeneAmp PCR System 9700 (Perkin Elmer). The linear amplification started with a 10 s denaturation at  $96^{\circ}$ C, 5 s annealing at  $50^{\circ}$ C and 4 min extension at 60°C. The fluorescently labeled sequencing products were separated from residual reaction reagents using a Centri-Sep spin column (Princeton Separations, Aldelphia NJ) and electrophoresed on POP6 capillary at 1500 V for 30 min. Sequencing data were automatically collected and analyzed by the ABI Prism 310 software.

# Table A: Primers Used to Determine Genomic Organization of Cathepsin C

Region	Primer Sequences
Intron 1	F: 5'- TGTCAACTGCTCGGTTATGGGTAA-3' (#6) R: 5'- TCGAGCTTCTCTTCGTACACCACT-3' (#7)
Intron 2	F: 5'- TGACTACAAGTGGTTTGCCTTTTT-3' (#8) R: 5' TGCTGCCCTCTTCTTTATACTGC-3' (#9)
Intron 3	F: 5'- GCCTCTGAGAATGTGTATGTCAAC-3' (#10) R: 5' CCTGCCCCAAAAATGAGATA-3" (#11)

PCT/US00/20400 WO 01/07663

maon .	F: 5' TCGAAAAATCCCAAGGTAATC-3' R: 5' GGGCCTAGAAAGGAAATATACATT-3	(#12) ' (#13)
Intron 5	F: 5' AATTTGTTCGGAACTATTTATTGA-3' R: 5' TCGCTTCTAGCATACCCATA-3'	(#14) (#15)

Exon	Primer Sequences	
1	F: 5'-GGCGATCAGACTGGCACACTAGAA -3'	(#16)
•	R: 5'-CTTACCCATAACCGAGCAGTTGAC-3'	(#17)
2	F: 5'- GCAGACTGTGCTCAAACTGGGTAG-3'	(#18)
<b>4</b>	R: 5'-TCTACTAATCAGAAGAGGTTTCAG-3'	(#19)
3	F: 5'-GGCACATTTACTGTGAATGAGAG-3'	(#20)
	R: 5'-GTCTCATTTGTAGCAACTCAC-3'	(#21)
4	F: 5'-CCACTTTCCACTTAGGCACAG-3'	(#22)
	R: 5'-AGGATGGTATTCAGCATTCATA-3'	(#23)
5	F: 5'-ATCCTAGCTAGTCTGGTAGCTGAA-3'	(#24)
	R: 5'-TCTAGGTATCCCCGAAATCCATCA-3'	(#25)
6	F: 5'-GATTCTCTGTGAGGCTTCAGATGT-3'	(#26)
	R: 5'-GCCAACAACAGCCAGCTGCACACA-3'	(#27)
7	F: 5'-TCCCCACTTAACCACTTTGC-3'	(#28)
	R: 5'-ACTTCATAGCTGACCATCTTCC-3'	(#29)

#### Primers for cDNA templates:

5

10

15

20

30

F:5'-GCCGCCCTCCTGCTGCTTCT-3' (#30)

R: 5'-GGCTTAGGATTGGGGTCTGA-3'(#31)

We analyzed raw sequence data, generated consensus sequences, and produced nucleotide/amino acid alignments (DNASIS V2.6 for Windows, Hitachi Software Engineering Co., Ltd.). Mutations were detected by creating nucleotide/amino acid alignments of reported cathepsin C sequence data versus affected PLS patients sequence data using the Higgins-Sharpe UPGMA. Numbers in parentheses are SEQ ID NOS:.

#### Example I

#### PLS families

25

Parents of most families were consanguineous. Linkage studies localized a PLS gene in these five families to chromosome 11q14 [8]. Most affected individuals were homozygous for SSTR markers within the PLS candidate interval on chromosome 11q14, consistent with inheritance of both maternal and paternal copies of this genetic interval from a common familial ancestor ("identical by descent"). Four different haplotypes for short sequence tandem repeat (SSTR) markers spanning the 55

PCT/US00/20400

WO 01/07663

5

10

15

20

25

critical region were identified (Fig. 2), consistent with four independent mutations in the gene responsible for PLS.

#### Analysis of cathepsin C

Using RT-PCR, we found cathepsin C is normally expressed in epithelium from palms, soles, knees and keratinized oral gingiva from unaffected individuals (data not shown). The cathepsin C gene spans approximately 46 kb and consists of 7 exons. Sequence analysis of exonic, intronic and the 5'regulatory regions of the cathepsin C gene revealed PLS affected individuals from these families were homozygous for CTSC mutations that significantly altered the cathepsin C open reading frame.

Exon 6: Two affected individuals from one family were found to have an exon 6 nonsense mutation (856C->T) which introduces a stop codon at amino acid 286 (Fig. 3).

Exon 7: Three different exon 7 mutations were detected (Fig. 4). A deletion of a single nucleotide (1047delA) of codon 349 was found that introduced a frame shift and an early termination codon (TGA) 27 bases downstream. This mutation would result in a mutated protein of 358 amino acids, compared to the normal (wild type) 463 amino acids. A deletion of 2 bases of codon 343 (1028-1029delCT) resulting in the introduction of an early termination codon (TGA), and a truncated protein of 342 amino acids was identified in another family. A G->A substitution in codon 429 (1286G->A) that altered the original TRP codon (TGG) to a terminator codon (TAG) was identified in two affected individuals (#7 and #22) from two additional families. The expected truncated protein is 428 amino acids. Although these families were not known to be related, the fact that affected individuals from these two Turkish families are homozygous for a common cathepsin C

35

gene mutation and also share a common haplotype for SSTR markers in the PLS candidate interval flanking the cathepsin C gene (D11S931 - D11S1311) suggests that these individuals have inherited the same cathepsin C gene mutation from a common ancestor. Additional mutations were also identified in Exons 2, 3, 4, 5, 6 and 7. Summaries of the mutations identified to date are set forth in Table I and locations are shown in Figure 5.

Papillon Lefevre syndrome is a palmoplantar keratoderma (PPK) with the characteristic clinical features of palmoplantar hyperkeratosis and severe periodontal destruction. The PPKs are a heterogeneous group of diseases all having gross thickening of the palmoplantar skin. Clinically, the finding that distinguishes PLS from other PPKs is severe, early onset periodontal destruction. In affected individuals, the development and eruption of the primary teeth proceed normally, but the eruption of these teeth into the oral cavity is associated with gingival inflammation and subsequent rapid destruction of the periodontium. This form of destructive periodontitis is characteristically unresponsive to traditional periodontal treatment modalities, and consequently, the primary dentition is usually exfoliated prematurely. After exfoliation, the inflammation subsides, and the gingiva resumes a healthy appearance. However, with the eruption of the permanent dentition the process is usually repeated, resulting in the premature exfoliation of the permanent dentition, although the third molars are sometimes spared [5]. Destruction of the alveolar bone in PLS is usually severe, resulting in generalized atrophy of the alveolar ridges, further complicating dental therapy.

Because cathepsin C both localized to the refined PLS candidate interval on chromosome 11q14 and was normally expressed in epithelium from sites affected by PLS it was evaluated as a candidate gene for PLS.

35

5

10

15

20

25

5

10

15

20

25

30

35

PCT/US00/20400

Cathepsin C, or dipeptidyl aminopeptidase I (EC 3.4.14.1), is a lysosomal cysteine protease capable of removing dipeptides from the terminus of protein substrates, but at higher pH it also exhibits dipeptidyl transferase activity [10]. The cathepsin C gene spans approximately 46 kb and consists of 7 exons that encode a 463-amino acid polypeptide with predicted features of the papain family of cysteine proteases [11]. Unlike cathepsin B, H, L, and S, which are small monomeric enzymes, cathepsin C is a large (200 kD) oligomeric protein that consists of four identical subunits, each composed of three different polypeptide chains [12,13]. Expression of cathepsin C (CTSC) is tissue dependent [14]. CTSC is expressed in pituitary gland, spinal cord, aorta, left atrium, right atrium, left ventricle, right ventricle, inter ventricular septum, apex of heart, esophagus, stomach, duodenum, jejunum, ileum, ileocecum, appendix, ascending colon, transverse colon, descending colon, rectum, kidney, skeletal muscle, spleen, thymus, peripheral blood lymphocytes, lymph node, bone marrow, trachea, lung, placenta, bladder, uterus, testis, liver, pancreas, adrenal gland, thyroid gland, salivary gland, mammary gland, fetal heart, fetal kidney, fetal liver, fetal spleen, fetal thymus, and fetal lung. The CTSC message is also expressed at high levels in immune cells including polymorphonuclear leukocytes and alveolar macrophages and is also expressed at high levels in osteoclasts [11,16]. The pathologic clinical findings of the PLS affected individuals studied here involve severe inflammation and destruction of the gingiva as well as hyperkeratosis of the skin from palmar, plantar and knee sites. In unaffected individuals, cathepsin C is normally expressed in epithelial tissues from sites clinically affected by PLS.

Most parents of the PLS affected individuals in As a result, most PLS this study are consanguineous.

PCT/US00/20400 WO 01/07663

affected individuals in each family are homozygous for cathepsin C mutations inherited from a common ancestor. Yet parents and several siblings who are heterozygous carriers for cathepsin C mutations do not appear to show either the palmoplantar hyperkeratosis or severe early onset periodontitis characteristic of PLS. that the presence of one wild type cathepsin C gene is sufficient to prevent PPK and periodontal destruction in most patients. However, 1 mutation identified to date, (1047 A->G) appears to be associated with the presence of dermatological lesions. A consistent finding in the three linkage reports to date is the lack of a common haplotype among affected individuals from different families. The present report describes 20 different cathepsin C gene mutations associated with PLS. These findings suggest that the CTSC mutations responsible for PLS have arisen independently. Several mutations reported here result in the introduction of premature stop codons. While the W429X mutation encodes a protein shortened by only 35 amino acids, the introduced stop codon is 1 amino acid from the asparagine residue in the active site (Fig. 5). It is likely that such a mutation would cause a conformational alteration that may decrease or abolish activity. Additionally, we have also identified single nucleotide changes that result in missense amino acid changes in several additional PLS affected individuals from other populations, suggesting that structural alterations of cathepsin C may cause PLS.

In addition to the cardinal features of PLS, reports suggest some PLS patients have an increased This generalized susceptibility to infections [5]. increased susceptibility to infection may reflect the more deleterious effects of specific cathepsin C mutations, or may reflect the epigenetic effects of other gene loci. A variety of immunological findings have been reported in PLS affected individuals including

35

5

10

15

20

25

decreased monocyte chemotaxis, decreased neutrophil chemotaxis, impaired neutrophil phagocytosis, altered superoxide production, and decreased blastogenic response, but it has been difficult to extrapolate results of these studies. Consequently, the underlying pathogenesis of PLS has been poorly understood [17]. Immunological findings previously reported for affected individuals from the current families includes decreased PMN chemotaxis and elevated CD11b expression [18,19]. The pathologic clinical findings associated with PLS suggest that cathepsin C is functionally important in the structural growth and development of skin and in susceptibility to periodontal disease. As a lysosomal cysteine proteinase, cathepsin C is important in intracellular degradation of proteins and appears to be a central coordinator for activation of many serine proteinases in immune/inflammatory cells [11]. unknown if the profound periodontal disease susceptibility is a consequence of altered integrity of junctional epithelium surrounding the teeth. interesting that once teeth are exfoliated, and consequently the junctional epithelium is eliminated, the severe gingival inflammation resolves. A more complete understanding of the functional physiology of cathepsin C carries significant implications for understanding periodontal disease susceptibility. Identification of cathepsin C gene mutations in PLS raises the possibility of creating an animal model to study the development, treatment and prevention of hyperkeratosis and periodontitis.

Classification of the PPKs based upon histological findings, epidermolysis and localization of lesions within the skin (diffuse, linear or focal) has not been helpful in understanding the pathomechanism of disease [20]. Identification of mutations in specific genes has led to development of a revised nosology of these diseases in which PLS is grouped with the palmoplantar

35

30

5

10

15

20

ectodermal dysplasias [2]. In addition to providing insight into both normal as well abnormal epithelial growth and development, identification of mutations in cathepsin C associated with PLS will contribute to the overall nosology of the PPKs.

5

10

15

20

25

30

35

#### EXAMPLE II

# CTSC Mutation in Haim-Munk Syndrome

Of the many palmoplantar keratoderma (PPK) conditions, only Papillon Lefevre syndrome (PLS) and Haim Munk syndrome (HMS) are associated with premature periodontal destruction. Although both PLS and HMS share the cardinal features of PPK and severe periodontitis, a number of additional findings are reported in HMS including arachnodactyly, acroosteolysis, atrophic changes of the nails, and a radiographic deformity of the fingers. While PLS cases have been identified throughout the world, HMS has only been described among descendents of a religious isolate originally from Cochin, India. Parental consanguinity is a characteristic of many cases of both conditions. Although autosomal recessive transmission of PLS is evident, a more "complex" autosomal recessive pattern of inheritance with phenotypic influences from a closely linked modifying locus has been hypothesized for HMS. As set forth in Example I, mutations of the cathepsin C gene have been identified as the underlying genetic defect in PLS. To determine if a cathepsin C mutation is also responsible for HMS, we sequenced the gene in affected and unaffected individuals from families with HMS. Here we report identification a mutation of cathepsin C (exon 6, 857A->G) that changes a highly conserved amino acid in the cathpesin C peptide. This mutation segregates with HMS in four nuclear families. Additionally, the existence of a shared common haplotype for genetic loci flanking the cathepsin C gene suggests that affected individuals descended from the Cochin

"identical by descent" from a common ancestor. This finding supports simple autosomal recessive inheritance for HMS in these families. As described above, we also report a mutation of the same exon 6 CTSC codon (856C->T) in a Turkish family with classic PLS. These findings provide evidence that PLS and HMS are allelic variants of cathepsin C gene mutations.

In addition to congenital palmoplantar keratosis and progressive early onset periodontal destruction, other clinical findings shared by these individuals included recurrent pyogenic skin infections, acroosteolysis, atrophic changes of the nails, arachnodactyly, and a peculiar radiographic deformity of the fingers consisting of tapered pointed phalangeal ends and a clawlike volar curve (Figures 6A and 6B). Subsequently pes planus was reported to be associated with the syndrome [24]. This was the first reported association of these clinical findings, and the condition became known as Haim Munk syndrome, or keratosis palmoplantaris with periodontopathia and onychogryposis (HMS1; MIM245010)[22]. Although the palmoplantar findings and severe periodontitis were suggestive of the Papillon-Lefevre syndrome (PLS; MIM245000)[3], the association of other clinical features, particularly nail deformities and arachnodactyly, argued that HMS was a distinct disorder. In contrast to PLS, the skin manifestations in HMS were reported to be more severe and extensive. In addition to a marked palmoplantar keratosis (Figure 6C, 6D), affected individuals had scaly erythematous and circumscribed patches on the elbows, knees, forearms, shins and dorsum of the hands. While the periodontium in HMS was reported to be less severely affected than in PLS, gingival inflammation and alveolar bone destruction are present and severe (Figure 6E, 6F). In a subsequent genetic study of this extended family, Hacham-Zadeh and

35

30

5

10

15

20

coworkers [25] concluded that the syndrome might not behave as a simple autosomal recessive trait. Based upon their estimate of the disease allele frequency in this population (0.1), the absence of the condition in other kindreds of the Cochin isolate, and an inability to document consanguinity for many of the parents of affected individuals, they hypothesized that a "complex" autosomal recessive inheritance pattern with a closely linked dominant modifier locus may be responsible for the HMS phenotype.

#### HMS families

5

10

15

20

25

Pedigrees of the reported familial relationships for the Cochin descendents are shown in Figure 7A.

Descendents of the Cochin isolate studied include sibships 2, 3, 4 and 5 in the kindred pedigree originally described by Hacham-Zadeh and coworkers [25].

# HMS family genotyping results

All HMS affected individuals from the Cochin kindred were found to be homozygous for all three polymorphic DNA loci (D11S1887, D11S1780 and D11S1367) flanking the cathepsin C locus. Additionally, these individuals shared a common haplotype for these polymorphic markers. These findings are consistent with inheritance of both maternal and paternal copies of this genetic interval from a common familial ancestor ("identical by descent").

# Analysis of cathepsin C in HMS

The cathepsin C gene spans approximately 46 kb and consists of 7 exons. Sequence analysis of exonic, intronic and the 5 regulatory regions of the cathepsin C gene revealed that HMS affected individuals from the Cochin kindred were homozygous for a mutation in codon 286 of exon 6 (857A->G) which results in substitution of a conserved glutamine residue at position 286 by an

35

arginine: Q286R (Figure 8). This glutamine residue is normally completely conserved in wild type cathepsin C from at least five species (data not shown). This was the only sequence change different from the reported, highly conserved, wild type CTSC sequence (GenBank Accession No.: AC011088; SEQ ID NO: 1). All available parents of HMS affected individuals were found to be heterozygous for the mutated (857A->G) allele and the wild type allele. None of the parents or siblings heterozygous for the mutated (857A->G) allele and the wild type allele manifested clinically identifiable characteristics of PPK or had a history of severe, early onset periodontitis.

### Restriction Analysis

5

10

15

20

25

30

35

The Q286R mutation creates an AvaI restriction cleavage site. We utilized this newly created restriction site to develop a rapid test to screen for the Q286R mutation. After amplification of a 465bp fragment encompassing the 3' end of exon 6 using primers: Forward 5'-GTATGCTAGAAGCGAGAATCCGTAT-3' (SEQ ID NO: 32) and Reverse 5'-CCAATGCTAAAACTTGTTGAGACC-3' (SEQ ID NO: 33), the PCR products were purified using the Promega PCR kit according to the manufacturer's instructions. Purified products were eluted in 20 µl water. Approximately 5-10 µl of purified product was digested with 5U AvaI (New England Biolabs) in a total volume of 15  $\mu$ l for 1.5 hr at 37°C. Following digestion, the products were separated by electrophoresis through an 1.8 % agarose gel. Amplification of the wildtype sequence results in a 465 bp product that is not cleaved by AvaI. Amplification of the mutated (857A(G) sequence results in a 465 bp product that is cleaved by AvaI to yield products of 404 and 61 bp. Accordingly, individuals who are homozygous for the wildtype sequence exhibit a 465 bp band. Heterozygous individuals exhibit 3 bands: 465, 404, and

PCT/US00/20400

WO 01/07663

61 bp bands. Individuals who are homozygous for the Q286R mutation exhibit bands of 404 and 61 bp. Restriction analysis confirmed the sequencing results of all examined individuals (Figure 9).

5

#### EXAMPLE III

# Genetic Screening for PPK-Associated Mutations

The foregoing findings provide the basis for screening and diagnostic assays for assessing patients for the presence of mutations in the CTSC gene related to the pathological conditions described herein. A summary of the mutations in CTSC identified as associated with PPKs are set forth in Table 1.

15

Table 1. Phenotype correlations with CTSC mutations

Mutation <sup>a</sup>	Predicted Effect	Location	Phenotype <sup>b</sup>	# of Families
199-222del	Deletion of aa 67-74	Exon 2	PLS	1
199-222dei 145-446insATGT	Frameshift, termination at aa 157	Exon 3	PLS	1
458С®Т	Threonine 153 Isoleucine	Exon 3	PLS	2
622-623insC	Frameshift, termination at aa 223	Exon 4	PLS	1
704G®A	Tryptophan 235 Stop	Exon 5	PLS	1
748C@T	Arginine 250 Stop	Exon 5	PLS	1
815G®C	Arginine 272 Proline	Exon 6	PLS	1
856C®T	Glutamine 286 Stop	Exon 6	PLS	2
857A®G	Glutamine 286 Arginine	Exon 6	HMS	1
898G®A	Glycine 300 Serine	Exon 7	PLS	1
901G®T	Glycine 301 Valine	Exon 7	PLS	1
901G®A	Glycine 301 Serine	Exon 7	PLS	1
910T®A	Tyrosine 304 Asparagine	Exon 7	PLS, RP	1
956A@G	Glutamic Acid 319 Glycine	Exon 7	PLS	1
1015C®T	Arginine 339 Cysteine	Exon 7	PLS	2
1019A©G	Tyrosine 340 Cysteine	Exon 7	PLS	1
1028-1029delCT	Introduction of premature termination codon	Exon 7	PLS	1
1040A®G	Tyrosine 347 Cysteine	Exon 7	PPP	2
1047delA	Frameshift, termination at aa 349	Exon 7	PLS	1
1286G®A	Tryptophan 429 Stop	Exon 7	PLS	3
1360A®G	Glutamic Acid 447 Glycine	Exon 7	PLS	1

- 1. cDNA numbering considering the initiator Met codon as nucleotide +1.
- 2. Phenotype symbols: PLS, Papillon-Lefevre syndrome; PPP, Prepubertal periodontitis;

HMS, Haim Munk syndrome; RP, Retinitis pigmentosa

while the mutations described in the previous examples are associated with certain pathological conditions, it is important to note that the CTSC gene contains many polymorphisms. Many of these genetic changes are not associated with the disease state. The genetic changes assessed by the methods of the present invention must be associated with the production of an aberrant CTSC protein. Accordingly, a suitable assay for diagnosing this disorder includes the step of differentiating harmless polymorphisms from those

25

30

35

5

10

15

20

mutations which give rise to PPKs and periodontal disorders. These include changes in the coding sequence which give rise to decreased mRNA stability as compared to wild type CTSC mRNA. Alternatively cathepsin C enzymatic activity can be compared between altered CTSC coding sequences and nucleic acids encoding the wild type enzyme. Such assays are well known in the art and need not be set forth here. See for example, McGuire et al., Archives of Biochemistry and Biophysics 295:280-8, 1992; McDonald et al., J. of Biological Chemistry 244:2693-26709, 1969; Metroione et al, Biochemistry 5:1597-1604, 1966; and Vanha-Perttula et al., Histochemie 5:170-181, 1965.

#### References:

5

10

15

20

25

30

35

40

- 1. Papillon, M.M., and Lefevre, P. Deux cas de keratodermie palmaire et plantaire symetrique familiale (maladie de Meleda) chez le frere et la soeur. Coexistence dans les deus cas d'alterations dentaires grabes. Bull Soc Fr Dermatol Syphilis 1924;31:82-87.
- 2. Stevens, H.P., Kelsell, D.P., Bryant, S.P., Bishop, D.T., Spurr, N.K., Weissenbach, J., Marger, D., Marger, R.S., Leigh, I.M. Linkage of an American pedigree with palmoplantar keratoderma and malignancy (palmoplantar ectodermal dysplasia type III) to 17q24. Literature survey and proposed updated classification of the keratodermas. Arch Dermatol. 1996;132:(6) 640-51.
- 3. Online Mendelian Inheritance in Man, OMIM (TM). Johns Hopkins University, Baltimore, MD. MIM Number: {MIM 245000}: {8/20/98}: World Wide Web URL: http://www.ncbi.nlm.nih.gov/omim/
  - 4. Gorlin, RJ, Sedano H, and Anderson VE (1964), RJ. The syndrome of palmar-plantar hyperkeratosis and premature periodontal destruction of the teeth. J Pediatr 1964; **65**:895-908.
  - 5. Haneke, E. The Papillon-Lefevre syndrome: keratosis palmoplantaris with periodontopathy. Report of a case and review of the cases in the literature. *Hum Genet* 1979;**51**:(1) 1-35.
  - 6. Laass, M.W., Hennies, H.C., Preis, S., Stevens, H.P., Jung, M., Leigh, I.M., Wienker, T.F., Reis, A..

WO 01/07663

25

Localization of a gene for Papillon-Lefevre syndrome to chromosome 11q14-q21 by homozygosity mapping. Hum Genet  $1997; \mathbf{101}: (3) 376-82$ .

- 7. Fischer, J., Blanchet-Bardon, C., Prud'homme, J.F., Pavek, S., Steijlen, P.M., Dubertret, L., Weissenbach, J. Mapping of Papillon-Lefevre syndrome to the chromosome 11q14 region. *Eur J Hum Genet* 1997; 5:(3)156-60.
- 8. Hart, T.C., Bowden, D.W., Ghaffar, K.A., Wang, W., Cutler, C.W., Cebeci, I., Efeoglu, A., Firatli, E. Sublocalization of the Papillon-Lefevre syndrome locus on 11q14-q21. Amer J Med Genet 1998;79:(2) 134-139.
- 9. Hart TC, Bowden DW, Hart PS, Walker SJ, Callison SA, Bobby PL, Firatli E. An integrated physical and genetic map of the PLS locus interval on chromosome 11g14. Mammalian Genome In Press
- 10.Kirschke, H., Barrett, A.J., Rawlings, N.D.
  Proteinases 1: Lysosomal cysteine proteinases. In:
  Protein Profile Vol.2, P. Sheterline, ed. (London, UK:
  Academic Press Ltd.) 1995;1587-1643.
  - 11. Rao, N.V., Rao, G.V., Hoidal, J.R. Human dipeptidylpeptidase I. *J Biol Chem*.1997; **272**:10260-10265.
- 12. Paris, A., Strukelj, B., Pungercar, J., Renko, M., Dolenc, I., Turk, V. Molecular cloning and sequence analysis of human preprocathepsin C. FEBS Lett 1995;369:326-330.
- 13. Dolenc, I., Turk, B., Pungercic, G., Ritonja, A., Turk, V. Oligomeric structure and substrate induced inhibition of human cathepsin C. *J Biol Chem* 1995; **270**:21626-31.
- 14. Pham, C. T. N., Armstrong, R. J., Zimonjic, D.B., Popescu, N. C., Payan, D. G., Ley, T. J. Molecular cloning, chromosomal localization, and expression of murine dipeptidyl peptidase I. *J Biol Chem* 1997;272:10695-10703.
- 15. McGuire, M.J., Lipsky, P.E., Thiele, D.L. Cloning nad characterization of the cDNA encoding mouse dipeptidyl peptidase I (cathepsin C). Biochimica et Biophysica Acta 1997; 1351:267-73.
- 16. Hakeda, Y., Kumegawa, M. Osteoclasts in bone metabolism. Kaibogaku Zasshi *J Anat* 1991; **66**:215-225.
  - 17. Hart , T.C. and Shapira, L. Papillon Lefevre syndrome. Periodontol 2000 1994;6, 88-100.

18. Firatli, E., Tuzun, B., Efeoglu, A. Papillon-Lefevre syndrome. Analysis of neutrophil chemotaxis. *J Periodontol* 1996; **67**:617-20.

5

19. Firatli, E., Gurel, N., Efeoglu, A., Badur, S. Clinical and immunological findings in 2 siblings with Papillon-Lefevre syndrome. *J Periodontol* 1996; 67:1210-5.

10

20. Itin, P.H. Classification of autosomal dominant palmoplantar keratoderme: past-present-future. *Dermatol* 1992;**185**:163-165.

15

21. Haim, S., Munk J: Keratosis palmo-plantaris congenita, with periodontosis, arachnodactyly and peculiar deformity of the terminal phalanges. Br J Dermatol 1965;77:42-45.

20

22. Online Mendelian Inheritance in Man, OMIM (TM). Johns Hopkins University, Baltimore, MD. MIM Number: {MIM 245010}: {5/20/97}: World Wide Web URL: http://www.ncbi.nlm.nih.gov/omim/

25

23.Gorlin, R. J., Pindborg, J. J., Cohen, M. M., Jr. Syndromes of the Head and Neck. New York: McGraw-Hill (pub.) (2nd ed.) 1976. Pp. 373-376.

30

24. Puliyel JM, Sridharan Iyer KS. A syndrome of keratosis palmo-plantaris congenita, pes planus, onychogryphosis, periodontosis, arachnodactyly and a peculiar acro-osteolysis. Brit J Derm 1986;115:243-248.

35

25. Hacham-Zadeh S, Schaap T, Cohen MM. A genetic analysis of the Papillon-Lefevre syndrome in a Jewish family from Cochin. Am J Med Genet 1978;2:153-157.

26 Gorlin RJ, Cohen MM jr., Levin LS. Syndromes of the head and neck . 3rd edn. New York. Oxford University Press, 1990:853-854 (Oxford Monographs on Medical Genetics, No.19).

40

27. Hart T, Hart PS, Bowden D, Walker S, Callison S, Michalec M, Firatli E. Mutations of the cathepsin C gene are responsible for Papillon Lefevre syndrome. In Press J Med Genet

45

28. Hart TC, Stabholz A, Meyle J, Shapira L, Van Dyke TE, Cutler CW, Soskolne WA. Genetic studies of syndromes with severe periodontitis and palmoplantar hyperkeratosis. J Periodont Res 1997;32:81-89.

50

29. Weissenbach J, Gyapay G, Dib C, Vignal A, Morissette J, Millasseau P, Vaysseix G, Lathrop M. A second-generation linkage map of the human genome.

Nature 1992;359:794-801.

30. Lucker GP, Van de Kerkhof PC, Steijlen PM. The hereditary palmoplantar keratoses: an updated review and classification. Brit J Derm 1994;131(1):1-14.

- 31. Smith P, Rosenzweig KA. Seven cases of Papillon Lefevre Syndrome. Periodontics 1967;5:42-6.
- 32. Cohen T, Bloch N. Immigrant Jews from Cochin. In Goldschmidt E (ed) The Genetics of Migrant and Isolate Populations. Baltimore: Williams & Wilkins 1963;352.
- 33. Wilkie AO. Craniosynostosis: genes and mechanisms. Hum Molec Genet 1997;6:1647-56.
  - 34. Hola-Jamriska L, Tort JF, Dalton JP, Day SR, Fan J, Asskov J and Brindley PJ. Cathepsin C from Schistosoma japonicum; cDNA encoding the preproenzyme and its phylogenetic relationships. Eur J Biochem 1998;255:527-534.
  - 35. Wolters PJ, Raymond WW, Blount JL, Caughey GH. Regulated expression, processing, and secretion of dog mast cell dipeptidyl peptidase I. J Biol Chem 1998;25:15514-15520.
    - 36. Mackenzie IC, Rittman G, Gao Z, Leigh I, Lane EB. Patterns of cytokeratin expression in human gingival epithelia. J Periodont Res 1991;26468-478.
    - 37. Hormia M, Sahlberg C, Thesleff I, Airenne T. The epithelium-tooth interface—a basal lamina rich in laminin-5 and lacking other known laminin isoforms. J Den Res (1998);77:1479-85.
    - 38. Stabholz A, Taichman NS, Soskolne WA. Occurrence of Actinobacillus actinomycetemcomitans and anti leukotoxin antibodies in some members of an extended family affected by Papillon-Lefevre syndrome. J Periodontol 1995;66:653-57.
    - 39. Page RC. Altman LC. Ebersole JL. Vandesteen GE. Dahlberg WH. Williams BL. Osterberg SK. Rapidly progressive periodontitis. A distinct clinical condition. Journal of Periodontology. 54(4):197-209, 1983
  - 40. Page RC. Bowen T. Altman L. Vandesteen E. Ochs H.

    Mackenzie P. Osterberg S. Engel LD. Williams BL.

    Prepubertal periodontitis. I. Definition of clinical disease entity. Journal of Periodontology. 54(5):257-71, 1983

5

20

25

30

35

40

PCT/US00/20400

WO 01/07663

While certain of the preferred embodiments of the present invention have been described and specifically exemplified above, it is not intended that the invention be limited to such embodiments. Various modifications may be made thereto without departing from the scope and spirit of the present invention, as set forth in the following claims.

10